

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
14 June 2001 (14.06.2001)

PCT

(10) International Publication Number  
**WO 01/42285 A2**

(51) International Patent Classification<sup>7</sup>: **C07K 14/00**

(21) International Application Number: **PCT/US00/32990**

(22) International Filing Date: 5 December 2000 (05.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/172,852 10 December 1999 (10.12.1999) US  
60/172,354 16 December 1999 (16.12.1999) US

(71) Applicant (*for all designated States except US*): **INCYTE GENOMICS, INC.** [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **YUE, Henry** [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). **TANG, Y., Tom** [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). **LAL, Preeti** [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). **BURFORD, Neil** [GB/US]; 105 Wildwood Circle, Durham, CT 06422 (US). **AZIMZAI, Yalda** [US/US]; 5518 Boulder Canyon Drive, Castro Valley, CA 94552 (US). **PATTERSON, Chandra** [US/US]; 490 Sherwood Way, #1, Menlo Park, CA 94025 (US). **BAUGHN, Mariah, R.** [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). **LU, Dyung, Aina,**

**M.** [US/US]; 233 Coy Drive, San Jose, CA 95123 (US). **SHAH, Purvi** [IN/US]; 859 Salt Lake Drive, San Jose, CA 95133 (US). **AU-YOUNG, Janice** [US/US]; 233 Golden Eagle Lane, Brisbane, CA 94005 (US).

(74) Agents: **HAMLET-COX, Diana et al.**; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— *Without international search report and to be republished upon receipt of that report.*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **EXTRACELLULAR MATRIX AND CELL ADHESION MOLECULES**

(57) Abstract: The invention provides human extracellular matrix and cell adhesion molecules (XMAD) and polynucleotides which identify and encode XMAD. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of XMAD.

WO 01/42285 A2

## EXTRACELLULAR MATRIX AND CELL ADHESION MOLECULES

### TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of extracellular matrix and cell  
5 adhesion molecules and to the use of these sequences in the diagnosis, treatment, and prevention of  
genetic, autoimmune/inflammation, and cell proliferative disorders, including cancer, and in the  
assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid  
sequences of extracellular matrix and cell adhesion molecules.

### 10 BACKGROUND OF THE INVENTION

#### Extracellular Matrix Proteins

The extracellular matrix (ECM) is a complex network of glycoproteins, polysaccharides,  
proteoglycans, and other macromolecules that are secreted from the cell into the extracellular space.  
The ECM remains in close association with the cell surface and provides a supportive meshwork that  
15 profoundly influences cell shape, motility, strength, flexibility, and adhesion. In fact, adhesion of a cell  
to its surrounding matrix is required for cell survival except in the case of metastatic tumor cells, which  
have overcome the need for cell-ECM anchorage. This phenomenon suggests that the ECM plays a  
critical role in the molecular mechanisms of growth control and metastasis. (Reviewed in Ruoslahti, E.  
(1996) Sci. Am. 275:72-77.) Furthermore, the ECM determines the structure and physical properties  
20 of connective tissue and is particularly important for morphogenesis and other processes associated  
with embryonic development and pattern formation.

The collagens comprise a family of ECM proteins that provide structure to bone, teeth, skin,  
ligaments, tendons, cartilage, blood vessels, and basement membranes. Multiple collagen proteins have  
been identified. Three collagen molecules fold together in a triple helix stabilized by interchain disulfide  
25 bonds. Bundles of these triple helices then associate to form fibrils.

Elastin and related proteins confer elasticity to tissues such as skin, blood vessels, and lungs.  
Elastin is a highly hydrophobic protein of about 750 amino acids that is rich in proline and glycine  
residues. Elastin molecules are highly cross-linked, forming an extensive extracellular network of fibers  
and sheets. Elastin fibers are surrounded by a sheath of microfibrils which are composed of a number  
30 of glycoproteins, including fibrillin.

Fibronectin is a large ECM glycoprotein found in all vertebrates. Fibronectin exists as a dimer  
of two subunits, each containing about 2,500 amino acids. Each subunit folds into a rod-like structure  
containing multiple domains. The domains each contain multiple repeated modules, the most common  
of which is the type III fibronectin repeat. The type III fibronectin repeat is about 90 amino acids in

length and is also found in other ECM proteins and in some plasma membrane and cytoplasmic proteins. Furthermore, some type III fibronectin repeats contain a characteristic tripeptide consisting of Arginine-Glycine-Aspartic acid (RGD). The RGD sequence is recognized by the integrin family of cell surface receptors and is also found in other ECM proteins. (Reviewed in Alberts, supra, pp. 986-987.)

5 Laminin is a major glycoprotein component of the basal lamina which underlies and supports epithelial cell sheets. Laminin is one of the first ECM proteins synthesized in the developing embryo. Laminin is an 850 kilodalton protein composed of three polypeptide chains joined in the shape of a cross by disulfide bonds. Laminin is especially important for angiogenesis and, in particular, for guiding the formation of capillaries. (Reviewed in Alberts, B., et al. (1994) Molecular Biology of the  
10 Cell, Garland Publishing, New York, NY, pp. 990-991.)

Many proteinaceous ECM components are proteoglycans. Proteoglycans are composed of unbranched polysaccharide chains (glycosaminoglycans) attached to protein cores. Common proteoglycans include aggrecan, betaglycan, decorin, perlecan, serglycin, and syndecan-1. Some of these molecules not only provide mechanical support, but also bind to extracellular signaling molecules,  
15 such as fibroblast growth factor and transforming growth factor  $\beta$ , suggesting a role for proteoglycans in cell-cell communication. (Reviewed in Alberts, supra, pp. 973-978.)

Dentin phosphoryn (DPP) is a major component of the dentin ECM. DPP is a proteoglycan that is synthesized and expressed by odontoblasts (Gu, K., et al. (1998) *Eur. J. Oral Sci.* 106:1043-1047). DPP is believed to nucleate or modulate the formation of hydroxyapatite crystals.

20 Mucins are highly glycosylated glycoproteins that are the major structural component of the mucus gel. The physiological functions of mucins are cytoprotection, mechanical protection, maintenance of viscosity in secretions, and cellular recognition. MUC6 is a human gastric mucin that is also found in gall bladder, pancreas, seminal vesicles, and female reproductive tract (Toribara, N.W., et al. (1997) *J. Biol. Chem.* 272:16398-16403). The MUC6 gene has been mapped to human  
25 chromosome 11 (Toribara, N.W., et al. (1993) *J. Biol. Chem.* 268:5879-5885). Hemomucin is a novel Drosophila surface mucin that may be involved in the induction of antibacterial effector molecules (Theopold, U., et al. (1996) *J. Biol. Chem.* 271:12708-12715).

Extracellular matrix proteins may regulate cellular protein activity in a variety of ways.

Reversible protein phosphorylation is the primary method for regulating protein activity in eukaryotic  
30 cells. In general, proteins are activated by phosphorylation in response to extracellular signals such as hormones, neurotransmitters, and growth and differentiation factors. The activated proteins initiate the cell's intracellular response by way of intracellular signaling pathways and second messenger molecules such as cyclic nucleotides, calcium-calmodulin, inositol, and various mitogens, that regulate protein phosphorylation.

### Adhesion-Associated Proteins

The surface of a cell is rich in transmembrane proteoglycans, glycoproteins, glycolipids, and receptors. These macromolecules mediate adhesion with other cells and with components of the ECM. The interaction of the cell with its surroundings profoundly influences cell shape, strength, flexibility, motility, and adhesion. These dynamic properties are intimately associated with signal transduction pathways controlling cell proliferation and differentiation, tissue construction, and embryonic development.

Cadherins comprise a family of calcium-dependent glycoproteins that function in mediating cell-cell adhesion in virtually all solid tissues of multicellular organisms. These proteins share multiple repeats of a cadherin-specific motif, and the repeats form the folding units of the cadherin ECM. Cadherin molecules cooperate to form focal contacts, or adhesion plaques, between adjacent epithelial cells. The cadherin family includes the classical cadherins and protocadherins. Classical cadherins include the E-cadherin, N-cadherin, and P-cadherin subfamilies. E-cadherin is present on many types of epithelial cells and is especially important for embryonic development. P-cadherin is present on cells of the placenta and epidermis. Recent studies report that protocadherins are involved in a variety of cell-cell interactions (Suzuki, S. T. (1996) *J. Cell Sci.* 109:2609-2611). The intracellular anchorage of cadherins is regulated by their dynamic association with catenins, a family of cytoplasmic signal transduction proteins associated with the actin cytoskeleton. The anchorage of cadherins to the actin cytoskeleton appears to be regulated by protein tyrosine phosphorylation, and the cadherins are the target of phosphorylation-induced junctional disassembly (Aberle, H., et al. (1996) *J. Cell. Biochem.* 61:514-523).

Integrins are ubiquitous transmembrane adhesion molecules that link the ECM to the internal cytoskeleton. Integrins are composed of two noncovalently associated transmembrane glycoprotein subunits called  $\alpha$  and  $\beta$ . Integrins function as receptors that play a role in signal transduction. For example, binding of integrin to its extracellular ligand may stimulate changes in intracellular calcium levels or protein kinase activity (Sjaastad, M.D. and Nelson, W.J. (1997) *BioEssays* 19:47-55).

Lectins comprise a ubiquitous family of extracellular glycoproteins which bind cell surface carbohydrates specifically and reversibly, resulting in the agglutination of cells. (Reviewed in Drickamer, K. and Taylor, M.E. (1993) *Annu. Rev. Cell Biol.* 9:237-264.) This function is particularly important for activation of the immune response. Lectins mediate the agglutination and mitogenic stimulation of lymphocytes at sites of inflammation (Lasky, L.A. (1991) *J. Cell. Biochem.* 45:139-146; Palletta, E., et al. (1989) *J. Immunol.* 143:2850-2857). C-type lectin domains are found in a variety of proteins, including selectins and lecticans. Lecticans are a family of chondroitin sulfate proteoglycans that include aggrecan, versican, neurocan, and brevican. All C-type lectin proteins are



involved in protein-protein interactions (Aspberg, A., et al. (1997) Proc. Natl. Acad. Sci. USA 94:10116-10121). A novel macrophage-restricted C-type lectin protein has been cloned from mouse tissue. It is a type II transmembrane protein with one extracellular C-type lectin domain (Balch, S.G., et al. (1998) J. Biol. Chem. 273:18656-18664).

- 5           Toposome is a cell-adhesion glycoprotein isolated from mesenchyme-blastula embryos. Toposome precursors including vitellogenin promote cell adhesion of dissociated blastula cells.

LRRs are sequence motifs, approximately 22-28 amino acids in length, found in proteins with a large variety of functions and cellular locations. Proteins containing LRRs are all thought to be involved in protein-protein interactions. The crystal structure of LRRs has been studied and found to  
10   correspond to beta-alpha structural units. These structural units form a parallel beta sheet with one surface exposed to solvent. In this way an LRR-containing protein acquires a nonglobular shape (Kobe, B. and Deisenhofer, J. (1994) Trends Biochem. Sci. 19:415-421). There is evidence to suggest LRRs function in signal transduction and cellular adhesion as well as in protein-protein interactions (Gay, N.J., et al. (1991) FEBS Lett. 29:87-91).

- 15           Various proteins such as those encoded by the Drosophila armadillo gene and the human APC gene contain amino acid repeats that interact with  $\beta$ -catenins. The armadillo gene is required for pattern formation within the embryonic segments and imaginal discs and is highly conserved. It is 63% identical to a human protein, plakoglobin, which is involved in adhesive junctions joining epithelial and other cells (Peifer, M. and Wieschaus, E. (1990) Cell 63:1167-1176). APC gene mutations appear to  
20   initiate inherited forms of human colorectal cancer and sporadic forms of colorectal and gastric cancer (Rubinfeld, B., et al. (1993) Science 262:1731-1734). The fact that the protein encoded by APC interacts with catenin suggests a link between tumor initiation and cell adhesion (Su, L.K., et al. (1993) Science 262:1734-1737).

- SH3 is a 60-70 amino acid motif found in a variety of signal transduction and cytoskeletal  
25   proteins. The SH3 domain is involved in mediating protein-protein interactions. Evidence suggests that the SH3 domains recognize a family of related domains or proteins in a variety of different tissues and species. One novel SH3 domain-containing protein is the 52 kilodalton focal adhesion protein (FAP52 or p52). FAP52 is localized to focal adhesions, specialized membrane domains in cultured cells that mediate the attachment of cells to the growth substratum and ECM. Focal adhesions consist of  
30   structural proteins, integrins, regulatory molecules, and signaling molecules and are involved in cell signaling. FAP52 may form part of this multimolecular complex that comprises focal adhesion sites (Merilainen, J., et al. (1997) J. Biol. Chem. 272:23278-23284).

The discovery of new extracellular matrix and cell adhesion molecules and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the

diagnosis, prevention, and treatment of genetic, autoimmune/inflammation, and cell proliferative disorders, including cancer, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of extracellular matrix and cell adhesion molecules.

5

### SUMMARY OF THE INVENTION

The invention features purified polypeptides, extracellular matrix and cell adhesion molecules, referred to collectively as "XMAD" and individually as "XMAD-1," "XMAD-2," "XMAD-3," "XMAD-4," "XMAD-5," "XMAD-6," "XMAD-7," "XMAD-8," "XMAD-9," "XMAD-10," "XMAD-11," "XMAD-12," "XMAD-13," "XMAD-14," "XMAD-15," "XMAD-16," "XMAD-17," "XMAD-18," "XMAD-19," "XMAD-20," and "XMAD-21." In one aspect, the invention provides an isolated polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-21.

The invention further provides an isolated polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-21. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:22-42.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of an amino acid sequence selected from the group

consisting of SEQ ID NO:1-21. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide comprising an amino acid  
5 sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of an amino acid sequence selected from the group  
10 consisting of SEQ ID NO:1-21. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a  
15 polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of an amino acid  
20 sequence selected from the group consisting of SEQ ID NO:1-21.

The invention further provides an isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, c) a  
25 polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide sequence  
30 selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising

at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and  
5 optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of  
10 SEQ ID NO:22-42, b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or  
15 absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a composition comprising an effective amount of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a naturally occurring amino acid sequence  
20 having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and a pharmaceutically acceptable excipient. In one embodiment, the composition comprises an amino acid sequence selected from the group consisting of  
25 SEQ ID NO:1-21. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional XMAD, comprising administering to a patient in need of such treatment the composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an  
30 amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21. The method comprises a)

exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional XMAD, comprising administering to a patient in need of such treatment the composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional XMAD, comprising administering to a patient in need of such treatment the composition.

The invention further provides a method of screening for a compound that specifically binds to a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21. The method comprises a) combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

The invention further provides a method of screening for a compound that modulates the activity of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21. The

method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a  
5 change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:22-42, the method comprising a)  
10 exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

The invention further provides a method for assessing toxicity of a test compound, said method comprising a) treating a biological sample containing nucleic acids with the test compound; b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20  
15 contiguous nucleotides of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of i) a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, ii) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, iii) a polynucleotide sequence complementary to i), iv) a polynucleotide sequence complementary to ii),  
20 and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence selected from the group consisting of i) a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, ii) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a  
25 polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42, iii) a polynucleotide sequence complementary to i), iv) a polynucleotide sequence complementary to ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide comprises a fragment of a polynucleotide sequence selected from the group consisting of i)-v) above; c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization complex in the  
30 treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

#### BRIEF DESCRIPTION OF THE TABLES

35 Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs),

clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding XMAD.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods, algorithms, and searchable databases used for analysis of XMAD.

5 Table 3 shows selected fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones encoding XMAD were isolated.

10 Table 5 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

### DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood  
15 that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an,"  
20 and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings  
25 as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in  
30 connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

### DEFINITIONS

"XMAD" refers to the amino acid sequences of substantially purified XMAD obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and

human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of XMAD. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of XMAD either by directly interacting with XMAD or by acting on components of the biological pathway in which XMAD participates.

An "allelic variant" is an alternative form of the gene encoding XMAD. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding XMAD include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as XMAD or a polypeptide with at least one functional characteristic of XMAD. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding XMAD, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding XMAD. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent XMAD. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of XMAD is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to a sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known



in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of XMAD. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of XMAD either by directly interacting with XMAD or by acting on components of the biological pathway in which XMAD participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind XMAD polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic XMAD, or of any oligopeptide thereof,

to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

“Complementary” describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement,

5 3'-TCA-5'.

A “composition comprising a given polynucleotide sequence” and a “composition comprising a given amino acid sequence” refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution.

Compositions comprising polynucleotide sequences encoding XMAD or fragments of XMAD may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

“Consensus sequence” refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (Applied Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

“Conservative amino acid substitutions” are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as

25 conservative amino acid substitutions.

Original Residue	Conservative Substitution
Ala	Gly, Ser
Arg	His, Lys
Asn	Asp, Gln, His
30 Asp	Asn, Glu
Cys	Ala, Ser
Gln	Asn, Glu, His
Glu	Asp, Gln, His
Gly	Ala
35 His	Asn, Arg, Gln, Glu
Ile	Leu, Val
Leu	Ile, Val
Lys	Arg, Gln, Glu

5	Met Phe Ser Thr Trp Tyr Val	Leu, Ile His, Met, Leu, Trp, Tyr Cys, Thr Ser, Val Phe, Tyr His, Phe, Trp Ile, Leu, Thr
---	---	---

---

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

A "fragment" is a unique portion of XMAD or the polynucleotide encoding XMAD which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:22-42 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:22-42, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:22-42 is useful, for

example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:22-42 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:22-42 and the region of SEQ ID NO:22-42 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

5 A fragment of SEQ ID NO:1-21 is encoded by a fragment of SEQ ID NO:22-42. A fragment of SEQ ID NO:1-21 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-21. For example, a fragment of SEQ ID NO:1-21 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-21. The precise length of a fragment of SEQ ID NO:1-21 and the region of SEQ ID NO:1-21 to which the fragment  
10 corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A "full-length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A "full-length" polynucleotide sequence encodes a "full-length" polypeptide sequence.

15 "Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in  
20 the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular  
25 biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between  
30 aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at  
35 <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis

programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

*Matrix: BLOSUM62*

*Reward for match: 1*

*Penalty for mismatch: -2*

*Open Gap: 5 and Extension Gap: 2 penalties*

*Gap x drop-off: 50*

*Expect: 10*

*Word Size: 11*

*Filter: on*

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters

of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with  
 5 polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (Apr-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

10        *Matrix: BLOSUM62*  
           *Open Gap: 11 and Extension Gap: 1 penalties*  
           *Gap x drop-off: 50*  
           *Expect: 10*  
           *Word Size: 3*  
 15        *Filter: on*

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150  
 20 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for  
 25 chromosome replication, segregation and maintenance.

The term "humanized antibody" refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a  
 30 complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e.,

binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency; and therefore hybridization specificity. Permissive  
5 annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic  
10 strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating  $T_m$  and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al., 1989, Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

15 High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance,  
20 sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is  
25 strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g.,  $C_0t$  or  $R_0t$  analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g.,  
30 paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune

disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

5 An "immunogenic fragment" is a polypeptide or oligopeptide fragment of XMAD which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of XMAD which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

10 The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of XMAD. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of XMAD.

15 The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a  
20 functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which  
25 comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an XMAD may involve lipidation, glycosylation,  
30 phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of XMAD.

"Probe" refers to nucleic acid sequences encoding XMAD, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are  
35 isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical



labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes. "Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that

hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing  
5 primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence.

10 This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence.  
15 Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be use to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated  
20 regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription, translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent,  
25 chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose  
30 instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic acids encoding X<sub>MAD</sub>, or fragments thereof, or X<sub>MAD</sub> itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For  
5 example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free,  
10 preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters,  
15 chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

20 "Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection,  
25 electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to  
30 animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in  
35 vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The

transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants, and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook, J. et al. (1989), supra.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternative splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

## THE INVENTION

The invention is based on the discovery of new human extracellular matrix and cell adhesion molecules (XMAP), the polynucleotides encoding XMAP, and the use of these compositions for the diagnosis, treatment, or prevention of genetic, autoimmune/inflammation, and cell proliferative disorders, including cancer.

Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding XMAP. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide

and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each XMAD were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries.

- 5 In some cases, GenBank sequence identifiers are also shown in column 5. The Incyte clones and GenBank cDNA sequences, where indicated, in column 5 were used to assemble the consensus nucleotide sequence of each XMAD and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each  
10 polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows homologous sequences as identified by BLAST analysis along with relevant citations, all of which are expressly incorporated by reference herein in their entirety; and column 7 shows analytical methods and in some cases, searchable databases to which the analytical methods were applied. The  
15 methods of column 7 were used to characterize each polypeptide through sequence homology and protein motifs.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding XMAD. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are  
20 useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:22-42 and to distinguish between SEQ ID NO:22-42 and related polynucleotide sequences. The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express XMAD as a fraction of total tissues expressing XMAD. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing XMAD as a fraction of total  
25 tissues expressing XMAD. Column 5 lists the vectors used to subclone each cDNA library.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding XMAD were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

- 30 SEQ ID NO:18 maps to chromosome 22 within the interval from the P terminus to 19.5 centiMorgans.

The invention also encompasses XMAD variants. A preferred XMAD variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the XMAD amino acid sequence, and which contains at least one functional or structural

characteristic of XMAD.

The invention also encompasses polynucleotides which encode XMAD. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:22-42, which encodes XMAD. The polynucleotide sequences of  
5 SEQ ID NO:22-42, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil; and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding XMAD. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least  
10 about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding XMAD. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:22-42 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:22-42.  
15 Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of XMAD.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding XMAD, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the  
20 invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring XMAD, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode XMAD and its variants are generally capable of  
25 hybridizing to the nucleotide sequence of the naturally occurring XMAD under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding XMAD or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which  
30 particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding XMAD and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode XMAD and

XMAD derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding XMAD or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:22-42 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in

“Definitions.”

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Applied Biosystems, Foster City CA), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Applied Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Applied Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding XMAD may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.)

Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) *Nucleic Acids Res.* 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al.

(1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060).

5 Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a  
10 GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library  
15 does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-  
20 specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Applied Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments  
25 which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode XMAD may be cloned in recombinant DNA molecules that direct expression of XMAD, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent  
30 amino acid sequence may be produced and used to express XMAD.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter XMAD-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic



oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Crameri, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or improve the biological properties of XMAD, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding XMAD may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucleic Acids Symp. Ser. 7:215-223; Horn, T. et al. (1980) Nucleic Acids Symp. Ser. 7:225-232.) Alternatively, XMAD itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Applied Biosystems). Additionally, the amino acid sequence of XMAD, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) Methods Enzymol. 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, supra, pp. 28-53.)

In order to express a biologically active XMAD, the nucleotide sequences encoding XMAD or

derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding X<sub>MAD</sub>. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding X<sub>MAD</sub>. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding X<sub>MAD</sub> and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding X<sub>MAD</sub> and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding X<sub>MAD</sub>. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509; Bitter, G.A. et al. (1987) *Methods Enzymol.* 153:516-544; Scorer, C.A. et al. (1994) *Bio/Technology* 12:181-184; Engelhard, E.K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227; Sandig, V. et al. (1996) *Hum. Gene Ther.* 7:1937-1945; Takamatsu, N. (1987) *EMBO J.* 6:307-311; Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659; and Harrington,

J.J. et al. (1997) Nat. Genet. 15:345-355.) Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) Cancer Gen. Ther. 5(6):350-356; Yu, M. et al. (1993) Proc. Natl. Acad. Sci. USA 90(13):6340-6344; Buller, R.M. et al. (1985) Nature 317(6040):813-815; McGregor, D.P. et al. 5 (1994) Mol. Immunol. 31(3):219-226; and Verma, I.M. and N. Somia (1997) Nature 389:239-242.) The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding XMAD. For example, routine cloning, 10 subcloning, and propagation of polynucleotide sequences encoding XMAD can be achieved using a multifunctional *E. coli* vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding XMAD into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for *in vitro* 15 transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of XMAD are needed, e.g. for the production of antibodies, vectors which direct high level expression of XMAD may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

20 Yeast expression systems may be used for production of XMAD. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast *Saccharomyces cerevisiae* or *Pichia pastoris*. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, *supra*; 25 Bitter, *supra*; and Scorer, *supra*.)

Plant systems may also be used for expression of XMAD. Transcription of sequences encoding XMAD may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be 30 used. (See, e.g., Coruzzi, *supra*; Broglie, *supra*; and Winter, *supra*.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., *The McGraw Hill Yearbook of Science and Technology* (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases 35 where an adenovirus is used as an expression vector, sequences encoding XMAD may be ligated into an

adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses X<sub>2</sub> in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of X<sub>2</sub> in cell lines is preferred. For example, sequences encoding X<sub>2</sub> can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk*<sup>-</sup> and *aprt*<sup>-</sup> cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech),  $\beta$  glucuronidase and its substrate  $\beta$ -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding XMAD is inserted within a marker gene sequence, transformed cells containing sequences encoding XMAD can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding XMAD under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding XMAD and that express XMAD may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of XMAD using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on XMAD is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding XMAD include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding XMAD, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding XMAD may be cultured under

conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode XMAD may be designed to contain signal sequences which direct secretion of XMAD through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding XMAD may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric XMAD protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of XMAD activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the XMAD encoding sequence and the heterologous protein sequence, so that XMAD may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, *supra*, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled XMAD may be achieved *in vitro* using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or

SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, <sup>35</sup>S-methionine.

XMAD of the present invention or fragments thereof may be used to screen for compounds that specifically bind to XMAD. At least one and up to a plurality of test compounds may be screened for specific binding to XMAD. Examples of test compounds include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of XMAD, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, e.g., Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which XMAD binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express XMAD, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or E. coli. Cells expressing XMAD or cell membrane fractions which contain XMAD are then contacted with a test compound and binding, stimulation, or inhibition of activity of either XMAD or the compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with XMAD, either in solution or affixed to a solid support, and detecting the binding of XMAD to the compound. Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

XMAD of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of XMAD. Such compounds may include agonists, antagonists, or partial or inverse agonists. In one embodiment, an assay is performed under conditions permissive for XMAD activity, wherein XMAD is combined with at least one test compound, and the activity of XMAD in the presence of a test compound is compared with the activity of XMAD in the absence of the test compound. A change in the activity of XMAD in the presence of the test compound is indicative of a compound that modulates the activity of XMAD. Alternatively, a test compound is combined with an in vitro or cell-free system comprising XMAD under conditions suitable for XMAD activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of XMAD may do so indirectly and need not come in direct contact with the

test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding XMAD or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent No. 5,175,383 and U.S. Patent No. 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding XMAD may also be manipulated in vitro in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding XMAD can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding XMAD is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress XMAD, e.g., by secreting XMAD in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-74).

## THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of XMAD and extracellular matrix and cell adhesion molecules. In addition, the expression of XMAD is closely associated with cell proliferation. Therefore, XMAD appears to play



a role in genetic, autoimmune/inflammation, and cell proliferative disorders, including cancer. In the treatment of disorders associated with increased XMAD expression or activity, it is desirable to decrease the expression or activity of XMAD. In the treatment of disorders associated with decreased XMAD expression or activity, it is desirable to increase the expression or activity of XMAD.

5           Therefore, in one embodiment, XMAD or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of XMAD. Examples of such disorders include, but are not limited to, a genetic disorder, such as adrenoleukodystrophy, Alport's syndrome, choroideremia, Duchenne and Becker muscular dystrophy, Down's syndrome, cystic fibrosis, chronic granulomatous disease, Gaucher's disease, 10   Huntington's chorea, Marfan's syndrome, muscular dystrophy, myotonic dystrophy, pycnodysostosis, Refsum's syndrome, retinoblastoma, sickle cell anemia, thalassemia, Werner syndrome, von Willebrand's disease, Wilms' tumor, Zellweger syndrome, peroxisomal acyl-CoA oxidase deficiency, peroxisomal thiolase deficiency, peroxisomal bifunctional protein deficiency, mitochondrial carnitine palmitoyl transferase and carnitine deficiency, mitochondrial very-long-chain acyl-CoA 15   dehydrogenase deficiency, mitochondrial medium-chain acyl-CoA dehydrogenase deficiency, mitochondrial short-chain acyl-CoA dehydrogenase deficiency, mitochondrial electron transport flavoprotein and electron transport flavoprotein:ubiquinone oxidoreductase deficiency, mitochondrial trifunctional protein deficiency, and mitochondrial short-chain 3-hydroxyacyl-CoA dehydrogenase deficiency; an autoimmune/inflammation disorder, such as acquired immunodeficiency syndrome 20   (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, 25   erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, 30   ulcerative colitis, uveitis, Werner syndrome, complications of cancer; hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and a cell proliferative disorder, such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers 35   including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma,

and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus.

In another embodiment, a vector capable of expressing XMAD or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of XMAD including, but not limited to, those described above.

In a further embodiment, a composition comprising a substantially purified XMAD in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of XMAD including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of XMAD may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of XMAD including, but not limited to, those listed above.

In a further embodiment, an antagonist of XMAD may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of XMAD. Examples of such disorders include, but are not limited to, those genetic, autoimmune/inflammation, and cell proliferative disorders, including cancer, described above. In one aspect, an antibody which specifically binds XMAD may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express XMAD.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding XMAD may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of XMAD including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of XMAD may be produced using methods which are generally known in the art. In particular, purified XMAD may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind XMAD. Antibodies to XMAD may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit

dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with XMAD or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to XMAD have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of XMAD amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to XMAD may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda, S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce XMAD-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for XMAD may also be generated.

For example, such fragments include, but are not limited to,  $F(ab')_2$  fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the  $F(ab')_2$  fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al.

5 (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between XMAD and its

10 specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering XMAD epitopes is generally used, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for XMAD. Affinity is expressed as an association  
15 constant,  $K_a$ , which is defined as the molar concentration of XMAD-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The  $K_a$  determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple XMAD epitopes, represents the average affinity, or avidity, of the antibodies for XMAD. The  $K_a$   
20 determined for a preparation of monoclonal antibodies, which are monospecific for a particular XMAD epitope, represents a true measure of affinity. High-affinity antibody preparations with  $K_a$  ranging from about  $10^9$  to  $10^{12}$  L/mole are preferred for use in immunoassays in which the XMAD-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with  $K_a$  ranging from about  $10^6$  to  $10^7$  L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of XMAD, preferably in active form, from the antibody (Catty, D.

25 (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg  
30 specific antibody/ml, is generally employed in procedures requiring precipitation of XMAD-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al., supra.)

In another embodiment of the invention, the polynucleotides encoding XMAD, or any fragment

or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding XMAD. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding XMAD. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc., Totawa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, supra; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.* 25(14):2730-2736.)

In another embodiment of the invention, polynucleotides encoding XMAD may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) *Science* 270:404-410; Verma, I.M. and N. Somia (1997) *Nature* 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) *Nature* 335:395-396; Poeschla, E. et al. (1996) *Proc. Natl. Acad. Sci. USA.* 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the case where a genetic deficiency in XMAD expression or regulation causes disease, the expression of

XMAD from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in XMAD are treated by constructing mammalian expression vectors encoding XMAD and introducing these vectors by mechanical means into XMAD-deficient cells. Mechanical transfer technologies for use with cells *in vivo* or *ex vitro* include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) *Annu. Rev. Biochem.* 62:191-217; Ivics, Z. (1997) *Cell* 91:501-510; Boulay, J-L. and H. Récipon (1998) *Curr. Opin. Biotechnol.* 9:445-450).

Expression vectors that may be effective for the expression of XMAD include, but are not limited to, the PCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). XMAD may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or  $\beta$ -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Gossen, M. et al. (1995) *Science* 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) *Curr. Opin. Biotechnol.* 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and H.M. Blau, *supra*), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding XMAD from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al. (1982) *EMBO J.* 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to XMAD expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding XMAD under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive

element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) Proc. Natl. Acad. Sci. USA 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) J. Virol. 61:1647-1650; Bender, M.A. et al. (1987) J. Virol. 61:1639-1646; Adam, M.A. and A.D. Miller (1988) J. Virol. 62:3802-3806; Dull, T. et al. (1998) J. Virol. 72:8463-8471; Zufferey, R. et al. (1998) J. Virol. 72:9873-9880). U.S. Patent Number 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference.

Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4<sup>+</sup> T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) J. Virol. 71:7020-7029; Bauer, G. et al. (1997) Blood 89:2259-2267; Bonyhadi, M.L. (1997) J. Virol. 71:4707-4716; Ranga, U. et al. (1998) Proc. Natl. Acad. Sci. USA 95:1201-1206; Su, L. (1997) Blood 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding XMAD to cells which have one or more genetic abnormalities with respect to the expression of XMAD. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) Transplantation 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) Annu. Rev. Nutr. 19:511-544; and Verma, I.M. and N. Somia (1997) Nature 18:389:239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding XMAD to target cells which have one or more genetic abnormalities with respect to the expression of XMAD. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing XMAD to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) Exp. Eye Res. 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S.

Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) J. Virol. 73:519-532 and Xu, H. et al. (1994) Dev. Biol. 163:152-161, hereby incorporated by reference. The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding XMAD to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) Curr. Opin. Biotechnol. 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full-length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for XMAD into the alphavirus genome in place of the capsid-coding region results in the production of a large number of XMAD-coding RNAs and the synthesis of high levels of XMAD in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) Virology 228:74-83). The wide host range of alphaviruses will allow the introduction of XMAD into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have



been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

5        Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding XMAD.

10        Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of  
15        candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis.

20        Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding XMAD. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

25        RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and  
30        wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding XMAD.

Compounds which may be effective in altering expression of a specific polynucleotide may include,  
35        but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming

oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased XMAD expression or activity, a compound which specifically inhibits expression of the polynucleotide encoding XMAD may be therapeutically useful, and in the treatment of disorders associated with decreased XMAD expression or activity, a compound which specifically promotes expression of the polynucleotide encoding XMAD may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method commonly known in the art, including chemical modification of a compound known to be effective in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a polynucleotide encoding XMAD is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an *in vitro* cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding XMAD are assayed by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the polynucleotide encoding XMAD. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide can be carried out, for example, using a *Schizosaccharomyces pombe* gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruice, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruice, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable for use *in vivo*, *in vitro*, and *ex vivo*. For *ex vivo* therapy, vectors may be introduced into stem cells taken

from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

5 Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a composition which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient.

10 Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of XMAD, antibodies to XMAD, and mimetics, agonists, antagonists, or inhibitors of XMAD.

The compositions utilized in this invention may be administered by any number of routes  
15 including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case  
20 of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle  
25 injection, and obviates the need for potentially toxic penetration enhancers.

Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

Specialized forms of compositions may be prepared for direct intracellular delivery of  
30 macromolecules comprising XMAD or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, XMAD or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et

al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example XMAD or fragments thereof, antibodies of XMAD, and agonists, antagonists or inhibitors of XMAD, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the  $ED_{50}$  (the dose therapeutically effective in 50% of the population) or  $LD_{50}$  (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the  $LD_{50}/ED_{50}$  ratio. Compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the  $ED_{50}$  with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1  $\mu\text{g}$  to 100,000  $\mu\text{g}$ , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

## DIAGNOSTICS

In another embodiment, antibodies which specifically bind XMAD may be used for the diagnosis of disorders characterized by expression of XMAD, or in assays to monitor patients being

treated with XMAD or agonists, antagonists, or inhibitors of XMAD. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for XMAD include methods which utilize the antibody and a label to detect XMAD in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and  
5 may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring XMAD, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of XMAD expression. Normal or standard values for XMAD expression are established by combining body fluids or cell extracts  
10 taken from normal mammalian subjects, for example, human subjects, with antibody to XMAD under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of XMAD expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

15 In another embodiment of the invention, the polynucleotides encoding XMAD may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of XMAD may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of  
20 XMAD, and to monitor regulation of XMAD levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding XMAD or closely related molecules may be used to identify nucleic acid sequences which encode XMAD. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a  
25 conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding XMAD, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the XMAD encoding sequences. The hybridization probes of the subject  
30 invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:22-42 or from genomic sequences including promoters, enhancers, and introns of the XMAD gene.

Means for producing specific hybridization probes for DNAs encoding XMAD include the cloning of polynucleotide sequences encoding XMAD or XMAD derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may

be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as  $^{32}\text{P}$  or  $^{35}\text{S}$ , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

5 Polynucleotide sequences encoding XMAP may be used for the diagnosis of disorders associated with expression of XMAP. Examples of such disorders include, but are not limited to, a genetic disorder, such as adrenoleukodystrophy, Alport's syndrome, choroideremia, Duchenne and Becker muscular dystrophy, Down's syndrome, cystic fibrosis, chronic granulomatous disease, Gaucher's disease, Huntington's chorea, Marfan's syndrome, muscular dystrophy, myotonic  
10 dystrophy, pycnodysostosis, Refsum's syndrome, retinoblastoma, sickle cell anemia, thalassemia, Werner syndrome, von Willebrand's disease, Wilms' tumor, Zellweger syndrome, peroxisomal acyl-CoA oxidase deficiency, peroxisomal thiolase deficiency, peroxisomal bifunctional protein deficiency, mitochondrial carnitine palmitoyl transferase and carnitine deficiency, mitochondrial very-long-chain acyl-CoA dehydrogenase deficiency, mitochondrial medium-chain acyl-CoA  
15 dehydrogenase deficiency, mitochondrial short-chain acyl-CoA dehydrogenase deficiency, mitochondrial electron transport flavoprotein and electron transport flavoprotein:ubiquinone oxidoreductase deficiency, mitochondrial trifunctional protein deficiency, and mitochondrial short-chain 3-hydroxyacyl-CoA dehydrogenase deficiency; an autoimmune/inflammation disorder, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress  
20 syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis,  
25 glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner  
30 syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and a cell proliferative disorder, such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma,  
35 leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of

the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. The polynucleotide sequences encoding XMAD may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered XMAD expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding XMAD may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding XMAD may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding XMAD in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of XMAD, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding XMAD, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual

clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding XMAD may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding XMAD, or a fragment of a polynucleotide complementary to the polynucleotide encoding XMAD, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences encoding XMAD may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from the polynucleotide sequences encoding XMAD are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSSCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

Methods which may also be used to quantify the expression of XMAD include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid



quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described in Seilhamer, J.J. et al., "Comparative Gene Transcript Analysis," U.S. Patent No. 5,840,484, incorporated herein by reference. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

In another embodiment, antibodies specific for XMAP, or XMAP or fragments thereof may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by quantifying the number of expressed genes and their relative abundance under given conditions and at a given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent Number 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies, or other biological samples. The transcript image may thus reflect gene expression in vivo, as in the case of a tissue or biopsy sample, or in vitro, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with in vitro model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed

molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000) Toxicol. Lett. 112-113:467-471, expressly incorporated by reference herein). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties.

5 These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data  
10 after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at <http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is  
15 important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

In one embodiment, the toxicity of a test compound is assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the  
20 present invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

Another particular embodiment relates to the use of the polypeptide sequences of the present  
25 invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and  
30 analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, *supra*). The proteins are visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent  
35 such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is

generally proportional to the level of the protein in the sample. The optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of the present invention. In some cases, further sequence data may be obtained for definitive protein identification.

A proteomic profile may also be generated using antibodies specific for XMAD to quantify the levels of XMAD expression. In one embodiment, the antibodies are used as elements on a microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) *Anal. Biochem.* 270:103-111; Mendoze, L.G. et al. (1999) *Biotechniques* 27:778-788). Detection may be performed by a variety of methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J. Seilhamer (1997) *Electrophoresis* 18:533-537), so proteome toxicant signatures may be useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two

samples is indicative of a toxic response to the test compound in the treated sample.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. 5 (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.

In another embodiment of the invention, nucleic acid sequences encoding XMAP may be used 10 to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a 15 chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic 20 linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, e.g., Lander, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353-7357.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map 25 data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding XMAP on a physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as 30 linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized 35 by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences

mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

5           In another embodiment of the invention, XMAD, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between XMAD and the agent being tested may be measured.

10           Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with XMAD, or fragments thereof, and washed. Bound XMAD is then detected by methods well known in the art. Purified XMAD can  
15 also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

          In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding XMAD specifically compete with a test compound for binding XMAD.

20           In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with XMAD.

          In additional embodiments, the nucleotide sequences which encode XMAD may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such  
25 properties as the triplet genetic code and specific base pair interactions.

          Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

30           Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

          The disclosures of all patents, applications, and publications mentioned above and below, in  
35 particular U.S. Ser. No. 60/172,354, and U.S. Ser. No. 60/172,852, are hereby expressly incorporated

by reference.

## EXAMPLES

### 5 I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl  
10 cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A+) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN,  
15 Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP  
20 vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, *supra*, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000  
25 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), pcDNA2.1 plasmid (Invitrogen, Carlsbad CA), or pINCY plasmid (Incyte Genomics, Palo Alto CA). Recombinant  
30 plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 $\alpha$ , DH10B, or ElectroMAX DH10B from Life Technologies.

### II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by *in vivo* excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least

one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

### III. Sequencing and Analysis

Incute cDNA recovered in plasmids as described in Example II were sequenced as follows. Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Applied Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Applied Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, supra, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VI.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions, references, and threshold parameters. The first column of Table 5 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between two sequences). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software

Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

5       The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and PFAM to acquire annotation  
10       using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the  
15       GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

      The programs described above for the assembly and analysis of full length polynucleotide and  
20       amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:22-42. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

#### IV. Analysis of Polynucleotide Expression

      Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene  
25       and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel, 1995, supra, ch. 4 and 16.)

      Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is  
30       much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum \{length(Seq. 1), length(Seq. 2)\}}}$$



The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding XMAD occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table 3.

#### V. Chromosomal Mapping of XMAD Encoding Polynucleotides

The cDNA sequences which were used to assemble SEQ ID NO:22-42 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:22-42 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 5). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO:, to that map location.

The genetic map location of SEQ ID NO:18 is described in The Invention as a range, or interval, of a human chromosome. The map position of an interval, in centiMorgans, is measured

relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Génethon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Human genome maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

#### VI. Extension of XMAD Encoding Polynucleotides

The full length nucleic acid sequences of SEQ ID NO:22-42 were produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing  $Mg^{2+}$ ,  $(NH_4)_2SO_4$ , and  $\beta$ -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100  $\mu$ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5  $\mu$ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the

concentration of DNA. A 5  $\mu$ l to 10  $\mu$ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).

In like manner, the polynucleotide sequences of SEQ ID NO:22-42 are used to obtain 5' regulatory sequences using the procedure above, along with oligonucleotides designed for such extension, and an appropriate genomic library.

## **VII. Labeling and Use of Individual Hybridization Probes**

Hybridization probes derived from SEQ ID NO:22-42 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250  $\mu$ Ci of [ $\gamma$ -<sup>32</sup>P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10<sup>7</sup> counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human

genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

### VIII. Microarrays

The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, supra), mechanical microspotting technologies, and derivatives thereof. The substrate in each of the aforementioned technologies should be uniform and solid with a non-porous surface (Skena (1999), supra). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Skena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645; Marshall, A. and J. Hodgson (1998) Nat. Biotechnol. 16:27-31.)

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection. After hybridization, nonhybridized nucleotides from the biological sample are removed, and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

#### Tissue or Cell Sample Preparation

Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and poly(A)<sup>+</sup> RNA is purified using the oligo-(dT) cellulose method. Each poly(A)<sup>+</sup> RNA sample is

reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/ $\mu$ l oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/ $\mu$ l RNase inhibitor, 500  $\mu$ M dATP, 500  $\mu$ M dGTP, 500  $\mu$ M dTTP, 40  $\mu$ M dCTP, 40  $\mu$ M dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)<sup>+</sup> RNA with

5 GEMBRIGHT kits (Incyte). Specific control poly(A)<sup>+</sup> RNAs are synthesized by in vitro transcription from non-coding yeast genomic DNA. After incubation at 37°C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85°C to stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc.

10 (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14  $\mu$ l 5X SSC/0.2% SDS.

#### Microarray Preparation

15 Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5  $\mu$ g. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia

20 Biotech).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and

25 coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in US Patent No. 5,807,522, incorporated herein by reference. 1  $\mu$ l of the array element DNA, at an average concentration of 100 ng/ $\mu$ l, is loaded into the open capillary printing element by a high-speed robotic

30 apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60°C followed by washes in

35 0.2% SDS and distilled water as before.

### Hybridization

Hybridization reactions contain 9 µl of sample mixture consisting of 0.2 µg each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65 °C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm<sup>2</sup> coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 µl of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60 °C. The arrays are washed for 10 min at 45 °C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45 °C in a second wash buffer (0.1X SSC), and dried.

### Detection

Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC computer. The digitized data are displayed as an image where the signal intensity is mapped using a

linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

5 A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

#### IX. Complementary Polynucleotides

10 Sequences complementary to the XMAD-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring XMAD. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of XMAD. To inhibit transcription, a  
15 complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the XMAD-encoding transcript.

#### X. Expression of XMAD

Expression and purification of XMAD is achieved using bacterial or virus-based expression  
20 systems. For expression of XMAD in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3).

25 Antibiotic resistant bacteria express XMAD upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of XMAD in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding XMAD by either homologous recombination or bacterial-mediated  
30 transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther.

35 7:1937-1945.)

In most expression systems, XMAD is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from XMAD at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified XMAD obtained by these methods can be used directly in the assays shown in Examples XI and XV.

#### **XI. Demonstration of XMAD Activity**

An assay for XMAD activity measures the disruption of cytoskeletal filament networks upon overexpression of XMAD in cultured cell lines. (Reznicek, G. A. et al. (1998) J. Cell Biol. 141:209-225.) cDNA encoding XMAD is subcloned into a mammalian expression vector that drives high levels of cDNA expression. This construct is transfected into cultured cells, such as rat kangaroo PtK2 or rat bladder carcinoma 804G cells. Actin filaments and intermediate filaments such as keratin and vimentin are visualized by immunofluorescence microscopy using antibodies and techniques well known in the art. The configuration and abundance of cytoskeletal filaments can be assessed and quantified using confocal imaging techniques. In particular, the bundling and collapse of cytoskeletal filament networks is indicative of XMAD activity.

Alternatively, an assay for XMAD activity measures the amount of cell aggregation induced by overexpression of XMAD. In this assay, cultured cells such as NIH3T3 are transfected with cDNA encoding XMAD contained within a suitable mammalian expression vector under control of a strong promoter. Cotransfection with cDNA encoding a fluorescent marker protein, such as Green Fluorescent Protein (Clontech), is useful for identifying stable transfectants. The amount of cell agglutination, or clumping, associated with transfected cells is compared with that associated with untransfected cells. The amount of cell agglutination is a direct measure of XMAD activity.

Alternatively, protein kinase activity is measured by quantifying the phosphorylation of a protein substrate by XMAD in the presence of gamma-labeled  $^{32}\text{P}$ -ATP. XMAD is incubated with the protein substrate,  $^{32}\text{P}$ -ATP, and an appropriate kinase buffer. The  $^{32}\text{P}$  incorporated into the substrate is separated from free  $^{32}\text{P}$ -ATP by electrophoresis and the incorporated  $^{32}\text{P}$  is counted using a radioisotope counter. The amount of incorporated  $^{32}\text{P}$  is proportional to the activity of XMAD. A determination of



the specific amino acid residue phosphorylated is made by phosphoamino acid analysis of the hydrolyzed protein.

## XII. Functional Assays

XMAD function is assessed by expressing the sequences encoding XMAD at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT plasmid (Life Technologies) and pCR3.1 plasmid (Invitrogen), both of which contain the cytomegalovirus promoter. 5-10  $\mu$ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2  $\mu$ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of XMAD on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding XMAD and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding XMAD and other genes of interest can be analyzed by northern analysis or microarray techniques.

### **XIII. Production of XMAD Specific Antibodies**

XMAD substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the XMAD amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, *supra*, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Applied Biosystems) using Fmoc chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for anti-peptide and anti-XMAD activity by, for example, binding the peptide or XMAD to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

### **XIV. Purification of Naturally Occurring XMAD Using Specific Antibodies**

Naturally occurring or recombinant XMAD is substantially purified by immunoaffinity chromatography using antibodies specific for XMAD. An immunoaffinity column is constructed by covalently coupling anti-XMAD antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing XMAD are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of XMAD (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/XMAD binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and XMAD is collected.

### **XV. Identification of Molecules Which Interact with XMAD**

XMAD, or biologically active fragments thereof, are labeled with <sup>125</sup>I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled XMAD, washed, and any wells with labeled XMAD complex are assayed. Data obtained using different concentrations of XMAD are used to calculate values for the number, affinity, and association of XMAD with the candidate molecules.

Alternatively, molecules interacting with XMAD are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989, Nature 340:245-246), or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

- XMAD may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent No. 6,057,101).

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention.

- Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	22	1424691	BEPINON01	998379R1 (KIDNTUT01), 1424691H1 (BEPINON01), SXAEO2538V1
2	23	1450801	PENITUT01	046316H1 (CORNNOT01), 1450801CT1 (PENITUT01), 1450801H1 (PENITUT01), 1671961H1 (BLADNOT05)
3	24	1597872	BRAINOT14	814997R1 (OVRTUT01), 814997T1 (OVRTUT01), 1412857T6 (BRAINOT12), 1438406F1 (PANCNOT08), 1597872H1 (BRAINOT14), 1797683H1 (PROSTUT05), 3346671H1 (BRAITUT24)
4	25	1674661	BLADNOT05	1655227F6 (PROSTUT08), 1674661H1 (BLADNOT05), 1675239F6 (BLADNOT05), 1879940F6 (LEUKNOT03), 2121172F6 (BRSTNOT07), 2157008F6 (BRAINOT09), 2672389F6 (KIDNNOT19), 3270393H1 (BRAINOT20), 3387668H1 (LUNGTUT17), 3685486H1 (HEAANOT01), 4103531H1 (BRSTTUT17), 4850546H1 (TESTNOT10), 5028429H1 (COLCDIT01), 5661414H1 (BRAUNOT01)
5	26	1689337	PROSTUT10	542204R1 (OVARNOT02), 961047R2 (BRSTTUT03), 1623395T6 (BRAITUT13), 1663607F6 (BRSTNOT09), 1689337H1 (PROSTUT10), 2898863R6 (THYMNON04), 3507526H1 (CONCNOT01)
6	27	1746392	STOMTUT02	682990H1 (UTRSNOT02), 1663009F6 (BRSTNOT09), 1746392H1 (STOMTUT02), 1746392T6 (STOMTUT02), 2079257F6 (ISLTNOT01), 3099537F6 (PTHYNOT03), 3111943H1 (BRSTNOT17), 3391682H1 (LUNGTUT28), 4747243F6 (SMCRUNT01)
7	28	1825182	LSUBNOT03	983441H1 (TONGTUT01), 1825182F6 (LSUBNOT03), 1825182H1 (LSUBNOT03), 1825369F6 (LSUBNOT03), SAQB00477F1, SAQB00879F1, SAQB01310F1, SAQB00187F1, SAQA02142F1, SAQA00159F1
8	29	2155541	BRAINOT09	871127T1 (LUNGAST01), 1309342R1 (COLNFET02), 1544021T1 (PROSTUT04), 2155541H1 (BRAINOT09), 2155541X15F1 (BRAINOT09)

Table 1 (Cont.)

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
9	30	2215706	SINTFET03	570718H1 (MMLR3DT01), 756160R1 (BRAITUT02), 1511501F1 (LUNGNOT14), 2215706F6 (SINTFET03), 2215706H1 (SINTFET03), 2648753F6 (OVARNOT10), 2804428H1 (PENCNOT01), 3092672T6 (BRSTNOT19), 3597972H1 (FIBPNOT01), 3604953H1 (LUNGNOT30), 3879505H1 (SPLNNOT11), 4506625F6 (OVARTD01), 4708188H1 (BRAIFET02), 4985942H1 (LIVRTUT10), 5151981H1 (HEARFET03), 5644791H1 (UTRSTMR01), 5862219H1 (BRAYDIT01)
10	31	2347692	TESTTUT02	075856R1 (THPIPEB01), 370791R1 (LUNGNOT02), 1502478F1 (BRAITUT07), 2347692H1 (TESTTUT02), 2825041F6 (ADRETUT06)
11	32	2579048	KIDNTUT13	841019R1 (PROSTUT05), 1352253F1 (LATRTUT02), 1414589F6 (BRAINOT12), 1427648F1 (SINTBST01), 1996374R6 (BRSTTUT03), 2579048H1 (KIDNTUT13)
12	33	2604493	LUNGUT07	901679X18 (BRSTTUT03), 927970X54R1 (BRAINOT04), 1435427F6 (PANCNOT08), 1484806F6 (CORPNOT02), 1962694T6 (BRSTNOT04), 1990921F6 (CORPNOT02), 2279985R6 (PROSNON01), 2279985T6 (PROSNON01), 2294223T6 (BRAINON01), 2604493H1 (LUNGUT07), 2707717H1 (PONSAT01), 3421936H1 (UCMCNOT04), 4769752H1 (BRATNOT02), 4989101H1 (LIVRTUT10), SAEA01968R1
13	34	2787182	BRSTNOT13	916228H1 (BRSTNOT04), 1251624F6 (LUNGFET03), 1440454F6 (THYRNOT03), 1664091F6 (BRSTNOT09), 1812788X17C1 (PROSTUT12), 1812788X21C1 (PROSTUT12), 2787182H1 (BRSTNOT13), 2824324F6 (ADRETUT06), 2882479T6 (UTRSTUT05), 3833543H1 (PANCNOT17), 60147041D2, 60147042B6, 60147044D2, SXAE05916V1, SXAE02927V1, 91670173
14	35	3096668	CERVNOT03	2373962F6 (ISLTNOT01), 2373962T6 (ISLTNOT01), 2762988H1 (BRSTNOT12), 3096668F6 (CERVNOT03), 3096668H1 (CERVNOT03), 3096668T6 (CERVNOT03), SCGA06156V1, SCGA11275V1, SCGA07741V1

Table 1 (Cont.)

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
15	36	3143411	HNT2AZS07	540532T6 (LNODNOT02), 852710R1 (NGANNOT01), 860567R1 (BRAITUT03), 1402106F6 (LATRTUT02), 3143411H1 (HNT2AZS07), 3143411R6 (HNT2AZS07), 5135819H1 (OVARBIT04)
16	37	3170835	BRSTNOT18	3170835H1 (BRSTNOT18), 3171275F6 (BRSTNOT18)
17	38	3550808	SYNONOT01	00101F1 (U937NOT01), 1353706T1 (LATRTUT02), 1426227F1 (SINTBST01), 1804230F6 (SINTNOT13), 2361183T6 (LUNGFET05), 2606392H1 (LUNGTUT07), 3550808H1 (SYNONOT01), SBAA00101F1
18	39	3683905	HEAANOT01	833556H1 (PROSNOT07), 1494051H1 (PROSNON01), 3683905H1 (HEAANOT01), 5512558H1 (BRADDIR01), 5700822H1 (DRGCNOT01), g1267581
19	40	4062841	BRAINOT21	1863239H1 (PROSNOT19), 1863239T6 (PROSNOT19), 4062841H1 (BRAINOT21)
20	41	6394358	UTRENOT10	875733R6 (LUNGAST01), 1312637T6 (BLADTUT02), 2296386R6 (BRSTNOT05), 2296386T6 (BRSTNOT05), 6394358H1 (UTRENOT10)
21	42	2847752		g4126329.v113.gs_6.edit.5p 1-6416; g3449297 5802-10044 5547763H1 (TESTNOC01) 8322-8532; 3373379H1 (CONNTUT05) 8722-8984; 3331371H1 (BRAIFET01) 9378-9641; 5376974H1 (BRAXNOT01) 9634-9772; 4015537F6 (BRAXNOT01) 9709-10226; 5921447H1 (BRAIFET02) 9963-10235; 4700084F6 (BRALNOT01) 10218-10739; 670937H1 (CRBLNOT01) 10517-10780; 3788576H1 (BRAHNOT05) 10608-10904; 5929111H1 (BRAIFET02) 10791-11069; 3278762T6 (STOMFET02) 11029-11627; 2847752R6 (HNT2AZS07) 11181-11648

Table 2

Polyptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence	Homologous Sequence	Analytical Methods and Databases
1 (1424691)	222	S25 T88 S155 S179 T191 T207 S90 S96 S113 S119	N7 N53 N68	M1-G29: Signal peptide E173-E181: 5- hydroxytryptamine 5A (serotonin) receptor R213-D215: Rgd cell interaction motif	g7649266 Sterile-alpha motif and leucine zipper containing kinase AZK	MOTIFS BLAST_GENBANK SPSCAN BLIMPS-PRINTS
2 (1450801)	228	S177 S99	N25	F20-G211: Leucine Rich Repeat	g3786312 Extracellular matrix protein (Nishiu, J. et al. Genomics (1998) 52:378- 381)	MOTIFS BLAST_GENBANK HMMER PFAM BLIMPS-PRINTS
3 (1597872)	386	S88 S137 S229 S364 T370		G237-P247: Insulin- like growth factor G202-R341: Spliceosome- associated protein	g4033606 Extensin (Kieliszewski M.J., Lampert D.T. Plant J. (1994) 5:157- 172)	MOTIFS BLAST_GENBANK BLIMPS-BLOCKS BLAST-DOMO

Table 2 (Cont.)

Polyptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence	Homologous Sequence	Analytical Methods and Databases
4 (1674661)	833	S46 T276 T418 S34 T57 T229 T302 S382 T429 S505 S826 S200 S364 S480 T523 S555 T561 S685 T701 S742 Y249 Y345 Y736	N106 N121 N310 N419 N522 N564	M1-V23: Signal peptide F53-K481: Semaphorin domain	g1110599 Semaphorin homolog (Inagaki, S. et al. FEBS Lett (1995) 370:269-272)	MOTIFS BLAST_GENBANK SIGPEPT SPSCAN BLAST-PFAM
5 (1689337)	410	T358 S394 S139 S249 S17 S343 T385 S401	N3	M1-A16: Signal peptide P240-Q255: Prokaryotic molybdopterin oxidoreductase L55-A65: Prepro- orexin signature R349-D351: Rgd cell interaction motif	g3450883 Fibroin (Gosline, J.M. et al. J. Exp. Biol. (1999) 202:3295-3303)	MOTIFS BLAST_GENBANK SPSCAN BLIMPS-BLOCKS BLIMPS-PRINTS
6 (1746392)	360	S217 S255 T344 S37 S8 T28 S69 S113 T182 S188 S224 S242 S250		E17-P356: Neurofilament triplet H	g310200 proline-rich proteoglycan	MOTIFS BLAST_GENBANK BLAST-DOMO
7 (1825182)	377	S18 S41 S140 T267 S38 S62 S120 T343	N97 N128 N135 N146	F20-G211: Leucine Rich Repeat	g188864 mucin (Shimomura, T., Blood (1990) 75:2349-2356)	MOTIFS BLAST_GENBANK HMMER-PFAM BLIMPS-PRINTS



Table 2 (Cont.)

Polyptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence	Homologous Sequence	Analytical Methods and Databases
8 (2155541)	182	T116		M1-F22: Signal peptide T42-D62: Neutrophil cytosol factor D170-E178: 7-fold repeat proteins I motif R60-D62: Rgd cell interaction motif	g6164953 vacuolar sorting protein VPS29 [Mus musculus] Edgar, A.J. and Polak, J.M. (2000) Biochem. Biophys. Res. Commun. 277:622-630	MOTIFS BLAST_GENBANK SPSCAN BLIMPS-PRINTS BLIMPS-PFAM
9 (2215706)	513	S239 S325 T14 S51 S71 S110 S137 S171 S208 S213 T219 S263 S268 S349 S394 S403 S404 T96 T118 S149 S239 S365 T475 S479	N69 N89 N338 N429	S94-Q108: Adrenocorticotrophin receptor S328-S335: "Phage" integrase family R196-D198: Rgd cell interaction motif	g4322670 Dentin phosphoryn P = 7.6e-07	MOTIFS BLIMPS-PRINTS BLIMPS-PFAM
10 (2347692)	361	T81 T53 S158 S257 T333 S128	N8 N51 N306 N324	G242-M268: C1q domain proteins L146-P150: Laminin G domain protein G110-P288: Fibrillar collagen carboxyl-terminus	g1562534 csdp single- stranded DNA binding protein	MOTIFS BLAST_GENBANK BLIMPS-BLOCKS BLIMPS-PFAM BLAST-DOMO
11 (2579048)	327	S18 S199 T55 T72 S73 T285 S51 S140 S177 S262	N158	K41-T55: Histone H5 signature R213-D215: Rgd cell interaction motif		MOTIFS BLIMPS-PRINTS

Table 2 (Cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence	Homologous Sequence	Analytical Methods and Databases
12 (2604493)	1 110	S213 T795 S8 T17 S40 S64 S82 T118 S132 S151 S170 S192 S249 T317 T357 T476 S601 S642 S673 S674 S701 S731 T795 S1015 S1059 S1073 T84 S236 S276 T292 T309 S337 S457 T506 T744 S749 S813 T945 S976 T1078	N168 N472 N640 N671 N672 N691 N698 N729 N747 N851 N966	V532-V565, L577- F598, M609-L622: Armadillo/beta- catenin-like repeats S511-H1025: Mouse p120 protein R971-D973: Rgd cell interaction motif	g1702924 p0071 Catenin- related protein (Hatzfeld M., Nachtsheim C. J. Cell Sci. (1996) 109:2767-2778)	MOTIFS BLAST_GENBANK HMMER-PFAM BLIMPS-PFAM BLAST-PRODOM
13 (2787182)	386	S123 T210 S265 S283 S317 S326 T330 T338 T44 S79 S100 S109 T127 T142 T170 T214 S332	N14 N173	C57-Q72: Transmembrane motif M279-M363: osteonectin	g3876060 Weak similarity with nitrogen fixation regulator	MOTIFS BLAST_GENBANK HMMER PROFILES CAN
14 (3096668)	181	S114 S116 S118 S120 S122 T124 S154 T173 T13	N17	F16-E25: Alpha-type calcitonin signature	g3393011 Clumping factor B (Ni Eidhin D. et al. Mol. Microbiol. 1998 30:245- 257)	MOTIFS BLAST_GENBANK BLIMPS-PRINTS

Table 2 (Cont.)

Polyptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence	Homologous Sequence	Analytical Methods and Databases
15 (3143411)	374	S291 S75 S92 T206 T214 T298 T315 T23 T37 T50 S51 S262 T263 S306	N109 N304	M1-G24: Signal peptide N224-W247, V328- N348: Transmembrane motif Y46-C63, W163-C176: C-type lectin domain	g3790610 Layilin (Borowsky M.L., Hynes R.O. (1998) J. Cell Biol. 143:429-442)	MOTIFS BLAST_GENBANK SIGDEPT SPSCAN HMMER BLAST PFAM BLIMPS-PRINTS PROFILESAN BLAST-DOMO
16 (3170835)	102	S45 T57 T44	N33	M1-T19: Signal peptide A55-103: Insect cuticle protein	g2565394 Cuticle 12	MOTIFS BLAST_GENBANK SPSCAN BLAST-PFAM PROFILESAN BLIMPS-PRINTS BLAST-PRODOM BLAST-DOMO
17 (3550808)	510	S76 S88 S128 S150 T152 T308 T448 T461 S49 S56 S110 T138	N72 N136 N193 N253 N352 N411	M1-G20: Signal peptide L5-G23: Transmembrane motif G33-F46: Pheromone B alpha-1 receptor	g294502 Olfactomedin (Yokoe H., Anholt R.R. Proc. Natl. Acad. Sci. USA (1993) 90:4655-9)	MOTIFS BLAST_GENBANK SIGDEPT SPSCAN HMMER BLIMPS-PRINTS

Table 2 (Cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence	Homologous Sequence	Analytical Methods and Databases
18 (3683905)	185	S11 S57 S173 T135		M1-L170: von Willebrand factor domain score M2-F15, R37-F51, V103-G111 M1-R171: collagen glycoprotein precursor	g2654431 Type XII collagen	MOTIFS BLAST_GENBANK HMMER-PFAM BLIMPS-PRINTS BLAST-PRODOM BLAST-DOMO
19 (4062841)	207	T83 S201 T74 S166 T174 S190		V93-T174: PDZ domain (Also known as DHR or GLGF) P91-V171: SH3 domain	g3885828 Lin-7-A (Irie M. et al. Oncogene (1999) 18:2811-2817)	MOTIFS BLAST_GENBANK HMMER-PFAM BLIMPS-PRINTS BLIMPS-PFAM BLIMPS-PRODOM BLAST-PRODOM BLAST-DOMO
20 (6394358)	238	S2 S96 S100 S12 S26 T149 S200 T203	N160	L3-A25, T64-E110, E209-R218: 7-fold repeat proteins (clathrin) R136-D138: Rgd cell interaction motif		MOTIFS BLIMPS-PFAM

Table 2 (Cont.)

Polypeptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence	Homologous Sequence	Analytical Methods and Databases
21	3298	S93 T307 S2706 S29 S222 S240 T266 T403 T421 T440 T457 T533 T567 T569 S658 T744 T756 T778 T831 S865 S880 S882 T931 T944 S983 T985 S1030 T1089 T1223 T1246 T1304 S1347 S1366 T1433 T1465 T1488 S1578 S1630 S1685 T1841 S1903 T2184 S2359 S2368 T2482 S2513 S2663 S2847 T2863 S2878 S2880 S2885 S2897 T2964 T2979 S2980 S3053 S3281 S186 S194 T200 T250 S409 S497 T527 S545 S833 T852 T1020 S1180 S1230 T1275 S1659 S1660 S1737 S1829 S1976 T2066 S2113 T2216 S2322 S2355 S2549 S2552 T2709 S2819 S2822 S2830 S2919 S2956 T2964 T3273 Y399 Y449 Y842 Y2299	N631 N846 N1181 N1221 N1316 N1326 N1648 N1712 N1769 N2034 N2163 N2182 N2372 N2460 N2492 N2683 N2732 N2794 N3240	L1536-Y1553 L2525-A2545 I2669-L2689 S2712-A2730 L2741-N2761: Transmembrane Domains M1-E31: Signal peptide Y329-A423; Y437- L535 Y549-V641; F655- L746 Y760-T848; Y862- N952 Y965-Q1057; F1071- V1159 Y1178-I1265: Cadherin domains C1378-C1431; C1438- C1469 C1478-C1512; C1725- C1756 C1931-C1962; C1966- C2000: EGF domains F1542-D1704; F1792- E1920: Laminin G domains C2110-A2137; C2588- L2613: GPCR signature	g3449288 MEGF2 [Rattus norvegicus]	MOTIFS BLAST_GENBANK HMMER SPSCAN HMMER-PFAM BLIMPS-BLOCKS PROFILES-PRINTS BLIMPS-PRINTS

Table 3

Polynucleotide SEQ ID NO:	Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
22	1-517 852-905	Cardiovascular (0.200) Gastrointestinal (0.200) Urologic (0.222)	Cell proliferation(0.400) Cell proliferation/Cell line (0.200) Inflammation/Trauma(0.300)	pSPORT1
23	1-2387	Reproductive (0.397) Gastrointestinal (0.132) Musculoskeletal (0.118)	Cell proliferation(0.559) Inflammation/Trauma(0.176) Other (0.118)	pINCY
24	1-901 1456-1471	Reproductive (0.262) Gastrointestinal (0.179) Nervous (0.179)	Cell proliferation(0.536) Inflammation/Trauma(0.297) Cell proliferation/Cell line (0.190)	pINCY
25	1-1928 1776-3293	Reproductive (0.294) Gastrointestinal (0.157) Nervous (0.157)	Cell proliferation(0.471) Inflammation/Trauma(0.373) Cell proliferation/Cell line (0.118)	pINCY
26	1-821 1312-1324	Reproductive (0.351) Hematopoietic/Immune (0.135) Nervous (0.135)	Cell proliferation(0.486) Cell proliferation/Cell line (0.243) Inflammation/Trauma(0.351)	pINCY
27	1-626 1034-1324	Reproductive (0.227) Nervous (0.182) Gastrointestinal (0.159)	Cell proliferation(0.432) Cell proliferation/Cell line (0.205) Inflammation/Trauma(0.228)	pINCY
28	1-2429	Gastrointestinal (1.000)	Cell proliferation(1.000)	pINCY
29	1-50 591-985	Reproductive (0.197) Nervous (0.164) Hematopoietic/Immune (0.145)	Cell proliferation(0.441) Inflammation/Trauma(0.454) Cell proliferation/Cell line (0.191)	pINCY
30	1-285 813-930 1145-3381	Nervous (0.235) Reproductive (0.235) Gastrointestinal (0.126)	Cell proliferation(0.445) Cell proliferation/Cell line (0.202) Inflammation/Trauma(0.252)	pINCY
31	1-82 1728-1803	Reproductive (0.267) Nervous (0.233) Other (0.117)	Cell proliferation(0.383) Inflammation/Trauma(0.383) Cell proliferation/Cell line (0.217)	pINCY
32	1-430 752-964 1405-1515	Reproductive (0.263) Nervous (0.193) Cardiovascular (0.140)	Cell proliferation(0.544) Cell proliferation/Cell line (0.158) Inflammation/Trauma(0.246)	pINCY

Table 3 (Cont.)

Polynucleotide SEQ ID NO:	Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
33	1-570 860-1573 1859-2494 2864-4416	Nervous (0.383) Reproductive (0.210) Gastrointestinal (0.111)	Cell proliferation(0.395) Inflammation/Trauma(0.334) Cell proliferation/Cell line (0.198)	pINCY
34	1-30 190-234 889-4428	Reproductive (0.280) Nervous (0.195) Hematopoietic/Immune (0.146)	Cell proliferation(0.415) Inflammation/Trauma(0.366) Cell proliferation/Cell line (0.146)	pINCY
35	1-189 264-1907	Reproductive (0.333) Cardiovascular (0.167) Developmental (0.167)	Cell proliferation(0.333) Cell proliferation/Cell line (0.333) Inflammation/Trauma(0.375)	pINCY
36	1-773 1742-1839	Nervous (0.312) Reproductive (0.312) Gastrointestinal (0.125)	Cell proliferation(0.531) Inflammation/Trauma(0.250) Cell proliferation/Cell line (0.125)	pSPORT1
37	1-503	Reproductive (1.000)	Cell proliferation(1.000)	pINCY
38	1-167 449-946 1541-2154	Gastrointestinal (0.723) Reproductive (0.128) Urologic (0.085)	Cell proliferation(0.447) Inflammation/Trauma(0.489) Trauma (0.170)	pINCY
39	1-431 666-733	Reproductive (0.450) Nervous (0.250) Urologic (0.100)	Cell proliferation(0.750) Inflammation/Trauma(0.200) Trauma (0.100)	pINCY
40	1-48 301-453 634-665	Reproductive (0.300) Cardiovascular (0.200) Hematopoietic/Immune (0.200)	Cell proliferation(0.400) Cell proliferation/Cell line (0.300) Inflammation/Trauma(0.400)	pINCY
41	1-276 553-741 820-1235	Hematopoietic/Immune (0.349) Nervous (0.163) Gastrointestinal (0.140)	Inflammation/Trauma(0.512) Cell proliferation(0.349) Cell proliferation/Cell line (0.209)	pINCY

Table 4

Polynucleotide SEQ ID NO:	Library	Library Comment
22	BEPINOT01	This normalized bronchial epithelium library was constructed from 5.12 million independent clones from the BEPINOT01 library. RNA was made from a bronchial epithelium primary cell line derived from a 54-year-old Caucasian male. The normalization and hybridization conditions were adapted from Soares et al., PNAS (1994) 91:9228, using a longer (24-hour) reannealing hybridization period.
23	PENITUT01	Library was constructed using RNA isolated from tumor tissue removed from the penis of a 64-year-old Caucasian male during penile amputation. Pathology indicated a fungating invasive grade 4 squamous cell carcinoma involving the inner wall of the foreskin and extending onto the glans penis. Patient history included benign neoplasm of the large bowel, atherosclerotic coronary artery disease, angina pectoris, gout, and obesity. Family history included malignant pharyngeal neoplasm, chronic lymphocytic leukemia, and chronic liver disease.
24	BRAINOT14	Library was constructed using RNA isolated from brain tissue removed from the left frontal lobe of a 40-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology for the associated tumor tissue indicated grade 4 gemistocytic astrocytoma.
25	BLADNOT05	Library was constructed using RNA isolated from bladder tissue removed from a 60-year-old Caucasian male during a radical cystectomy, prostatectomy, and vasectomy. Pathology for the associated tumor tissue indicated grade 3 transitional cell carcinoma. Carcinoma in-situ was identified in the dome and trigone. Patient history included tobacco use.
26	PROSTUT10	Library was constructed using RNA isolated from prostatic tumor tissue removed from a 66-year-old Caucasian male during radical prostatectomy and regional lymph node excision. Pathology indicated an adenocarcinoma (Gleason grade 2+3). Adenofibromatous hyperplasia was also present. The patient presented with elevated prostate specific antigen (PSA). Family history included prostate cancer and secondary bone cancer.
27	STOMTUT02	Library was constructed using RNA isolated from stomach tumor tissue obtained from a 68-year-old Caucasian female during a partial gastrectomy. Pathology indicated a malignant lymphoma of diffuse large-cell type. Previous surgeries included cholecystectomy. Patient history included thalassemia. Family history included acute leukemia, malignant neoplasm of the esophagus, malignant stomach neoplasm, and atherosclerotic coronary artery disease.



Table 4 (Cont.)

Polynucleotide SEQ ID NO:	Library	Library Comment
28	LSUBNOT03	Library was constructed using RNA isolated from submandibular gland tissue obtained from a 68-year-old Caucasian male during a sialoadenectomy. Family history included acute myocardial infarction, atherosclerotic coronary artery disease, and type II diabetes.
29	BRAINOT09	Library was constructed using RNA isolated from brain tissue removed from a Caucasian male fetus, who died at 23 weeks' gestation.
30	SINTFET03	Library was constructed using RNA isolated from small intestine tissue removed from a Caucasian female fetus, who died at 20 weeks' gestation.
31	TESTTUT02	Library was constructed using RNA isolated from testicular tumor removed from a 31-year-old Caucasian male during unilateral orchiectomy. Pathology indicated embryonal carcinoma.
32	KIDNTUT13	Library was constructed using RNA isolated from kidney tumor tissue removed from a 51-year-old Caucasian female during a nephroureterectomy. Pathology indicated a grade 3 renal cell carcinoma. Patient history included depressive disorder, hypoglycemia, and uterine endometriosis. Family history included calculus of the kidney, colon cancer, and type II diabetes.
33	LUNGTUT07	Library was constructed using RNA isolated from lung tumor tissue removed from the upper lobe of a 50-year-old Caucasian male during segmental lung resection. Pathology indicated an invasive grade 4 squamous cell adenocarcinoma. Patient history included tobacco use. Family history included skin cancer.
34	BRSTNOT13	Library was constructed using RNA isolated from breast tissue removed from the left medial lateral breast of a 36-year-old Caucasian female during bilateral simple mastectomy and total breast reconstruction. Pathology indicated benign breast tissue. Patient history included a breast neoplasm, depressive disorder, hyperlipidemia, chronic stomach ulcer, and an ectopic pregnancy. Family history included myocardial infarction, cerebrovascular disease, atherosclerotic coronary artery disease, hyperlipidemia, skin cancer, breast cancer, depressive disorder, esophageal cancer, bone cancer, Hodgkin's lymphoma, bladder cancer, and heart condition.
35	CERVNOT03	Library was constructed using RNA isolated from uterine cervical tissue removed from a 40-year-old Caucasian female during a vaginal hysterectomy with bilateral salpingo-oophorectomy and dilation and curettage. Pathology indicated secretory phase endometrium.

Table 4 (Cont.)

Polynucleotide SEQ ID NO:	Library	Library Comment
36	HNT2AZS07	This subtracted library was constructed from RNA isolated from an hNT2 cell line (derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor) treated for three days with 0.35 micromolar AZ. The hybridization probe for subtraction was derived from a similarly constructed library from untreated hNT2 cells. 3.08M clones from the AZ-treated library were subjected to three rounds of subtractive hybridization with 3.04M clones from the untreated library. Subtractive hybridization conditions were based on the methodologies of Swaroop et al. (NAR (1991) 19:1954) and Bonaldo et al. (Genome Research (1996) 6:791).
37	BRSTNOT18	Library was constructed using RNA isolated from diseased breast tissue removed from a 57-year-old Caucasian female during a unilateral simple extended mastectomy. Pathology indicated mildly proliferative breast disease. Patient history included breast cancer and osteoarthritis. Family history included type II diabetes, gallbladder and breast cancer, and chronic lymphocytic leukemia.
38	SYNONOT01	Library was constructed using RNA isolated from synovial tissue removed from a 75-year-old Caucasian male.
39	HEAANOT01	Library was constructed using RNA isolated from right coronary and right circumflex coronary artery tissue removed from the explanted heart of a 46-year-old Caucasian male during a heart transplantation. Patient history included myocardial infarction from total occlusion of the left anterior descending myocardial atherosclerotic coronary artery disease, hyperlipidemia, myocardial ischemia, dilated cardiomyopathy, left ventricular dysfunction, and tobacco abuse. Previous surgeries included cardiac catheterization. Family history included atherosclerotic coronary artery disease.
40	BRAINOT21	Library was constructed using RNA isolated from diseased brain tissue removed from the left frontal lobe of a 46-year-old Caucasian male during a lobectomy. Pathology indicated focal cortical and subcortical scarring of the left frontal lobe, characterized by cavitation and extensive reactive changes, including marked gliosis and hemosiderin deposition, consistent with a history of remote severe head trauma. GFAP was positive in astrocytes. The pattern of reactivity is that of reactive gliosis. Patient history included traumatic intracranial hemorrhage and brain injury with loss of consciousness following head trauma. Family history included cerebrovascular disease, cerebrovascular disease, and atherosclerotic coronary artery disease.

Table 4 (Cont.)

Polynucleotide SEQ ID NO:	Library	Library Comment
41	UTRENOT10	Library was constructed using polyA RNA isolated from pooled uterine endometrial tissue removed from three adult females during endometrial biopsy. Pathology indicated normal endometrium.
42		The Incyte cDNAs for SEQ ID NO:42 were derived from cDNA libraries constructed from brain, including tissues associated with Huntington's disease, Alzheimer's disease, and multiple sclerosis, as well as from pituitary, testicular, stomach, spinal cord, kidney, and prostate tissues, and from ovarian, cervical, pancreatic, and soft tissue tumors.

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25: 3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183: 63-98; and Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value= 1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value= 1.0E-8 or less Full Length sequences: fastx score= 100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S and J.G. Henikoff, Nucl. Acid Res., 19:6565-72, 1991. J.G. Henikoff and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37: 417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and, if applicable, Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol., 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25: 217-221.	Normalized quality score $\geq$ GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPSscan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12: 431-439.	Score=3.5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch et al. <u>supra</u> ; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

- 5           a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-21,  
            b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-21,  
            c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21, and  
10           d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-21.

2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1-21.

15           3. An isolated polynucleotide encoding a polypeptide of claim 1.

4. An isolated polynucleotide encoding a polypeptide of claim 2.

20           5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID NO:22-42.

6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.

25           7. A cell transformed with a recombinant polynucleotide of claim 6.

8. A transgenic organism comprising a recombinant polynucleotide of claim 6.

30           9. A method for producing a polypeptide of claim 1, the method comprising:  
            a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and

35           b) recovering the polypeptide so expressed.

10. An isolated antibody which specifically binds to a polypeptide of claim 1.

11. An isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of:

- a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42,
- b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:22-42,
- c) a polynucleotide sequence complementary to a),
- d) a polynucleotide sequence complementary to b), and
- e) an RNA equivalent of a)-d).

12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 11.

13. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and
- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.

15. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and
- b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

16. A composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

17. A composition of claim 16, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-21.

18. A method for treating a disease or condition associated with decreased expression of functional XMAD, comprising administering to a patient in need of such treatment the composition of claim 16.

19. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

20. A composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

21. A method for treating a disease or condition associated with decreased expression of functional XMAD, comprising administering to a patient in need of such treatment a composition of claim 20.

22. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

23. A composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.

24. A method for treating a disease or condition associated with overexpression of functional XMAD, comprising administering to a patient in need of such treatment a composition of claim 23.

25. A method of screening for a compound that specifically binds to the polypeptide of claim 1, said method comprising the steps of:

- a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and
- b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a



compound that specifically binds to the polypeptide of claim 1.

26. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, said method comprising:

- 5           a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,
- b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and
- c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound
- 10       with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

27. A method for screening a compound for effectiveness in altering expression of a target

15   polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,
- b) detecting altered expression of the target polynucleotide, and
- 20       c) comparing the expression of the target polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.

28. A method for assessing toxicity of a test compound, said method comprising:

- a) treating a biological sample containing nucleic acids with the test compound;
- 25       b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 11 or fragment thereof;
- 30       c) quantifying the amount of hybridization complex; and
- d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

<110> INCYTE GENOMICS, INC.  
 YUE, Henry  
 AZIMZAI, Yalda  
 TANG, Y. Tom  
 PATTERSON, Chandra  
 BAUGHN, Mariah R.  
 LU, Dyung Aina M.  
 SHAH, Purvi  
 LAL, Preeti  
 AU-YOUNG, Janice  
 BURFORD, Neil

<120> EXTRACELLULAR MATRIX AND CELL ADHESION MOLECULES

<130> PF-0760 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/172,852; 60/172,354

<151> 1999-12-10; 1999-12-16

<160> 42

<170> PERL Program

<210> 1

<211> 222

<212> PRT

<213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 1424691CD1

<400> 1  
 Met Arg Gln Ile Ala Ser Asn Thr Ser Leu Gln Arg Ser Gln Ser  
 1 5 10 15  
 Asn Pro Ile Leu Gly Ser Pro Phe Phe Ser His Phe Asp Gly Gln  
 20 25 30  
 Asp Ser Tyr Ala Ala Val Arg Arg Pro Gln Val Pro Ile Lys  
 35 40 45  
 Tyr Gln Gln Ile Thr Pro Val Asn Gln Ser Arg Ser Ser Ser Pro  
 50 55 60  
 Thr Gln Tyr Gly Leu Thr Lys Asn Phe Ser Ser Leu His Leu Asn  
 65 70 75  
 Ser Arg Asp Ser Gly Phe Ser Ser Gly Asn Thr Asp Thr Ser Ser  
 80 85 90  
 Glu Arg Gly Arg Tyr Ser Asp Arg Ser Arg Asn Lys Tyr Gly Arg  
 95 100 105  
 Gly Ser Ile Ser Leu Asn Ser Ser Pro Arg Gly Arg Tyr Ser Gly  
 110 115 120  
 Lys Ser Gln His Ser Thr Pro Ser Arg Gly Arg Tyr Pro Gly Lys  
 125 130 135  
 Phe Tyr Arg Val Ser Gln Ser Ala Leu Asn Pro His Gln Ser Pro  
 140 145 150  
 Asp Phe Lys Arg Ser Pro Arg Asp Leu His Gln Pro Asn Thr Ile  
 155 160 165  
 Pro Gly Met Pro Leu His Pro Glu Thr Asp Ser Arg Ala Ser Glu  
 170 175 180  
 Glu Asp Ser Lys Val Ser Glu Gly Gly Trp Thr Lys Val Glu Tyr  
 185 190 195  
 Arg Lys Lys Pro His Arg Pro Ser Pro Ala Lys Thr Asn Lys Glu  
 200 205 210  
 Arg Ala Arg Gly Asp His Arg Gly Trp Arg Asn Phe  
 215 220

<210> 2  
 <211> 228  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1450801CD1

<400> 2  
 Met Ile Leu His Asn Gln Ile Thr Gly Ile Gly Arg Glu Asp Phe  
 1 5 10 15  
 Ala Thr Thr Tyr Phe Leu Glu Glu Leu Asn Leu Ser Tyr Asn Arg  
 20 25 30  
 Ile Thr Ser Pro Gln Val His Arg Asp Ala Phe Arg Lys Leu Arg  
 35 40 45  
 Leu Leu Arg Ser Leu Asp Leu Ser Gly Asn Arg Leu His Thr Leu  
 50 55 60  
 Pro Pro Gly Leu Pro Arg Asn Val His Val Leu Lys Val Lys Arg  
 65 70 75  
 Asn Glu Leu Ala Ala Leu Ala Arg Gly Ala Leu Ala Gly Met Ala  
 80 85 90  
 Gln Leu Arg Glu Leu Tyr Leu Thr Ser Asn Arg Leu Arg Ser Arg  
 95 100 105  
 Ala Leu Gly Pro Arg Ala Trp Val Asp Leu Ala His Leu Gln Leu  
 110 115 120  
 Leu Asp Ile Ala Gly Asn Gln Leu Thr Glu Ile Pro Glu Gly Leu  
 125 130 135  
 Pro Glu Ser Leu Glu Tyr Leu Tyr Leu Gln Asn Asn Lys Ile Ser  
 140 145 150  
 Ala Val Pro Ala Asn Ala Phe Asp Ser Thr Pro Asn Leu Lys Gly  
 155 160 165  
 Ile Phe Leu Arg Phe Asn Lys Leu Ala Val Gly Ser Val Val Asp  
 170 175 180  
 Ser Ala Phe Arg Arg Leu Lys His Leu Gln Val Leu Asp Ile Glu  
 185 190 195  
 Gly Asn Leu Glu Phe Gly Asp Ile Ser Lys Asp Arg Gly Arg Leu  
 200 205 210  
 Gly Lys Glu Lys Glu Glu Glu Glu Glu Glu Glu Glu Glu  
 215 220 225  
 Glu Thr Arg

<210> 3  
 <211> 386  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1597872CD1

<400> 3  
 Met Val His Phe Gln Ala Ser Glu Val Gln Gln Leu Leu His Asn  
 1 5 10 15  
 Lys Phe Val Val Ile Leu Gly Asp Ser Ile Gln Arg Ala Val Tyr  
 20 25 30  
 Lys Asp Leu Val Leu Leu Leu Gln Lys Asp Ser Leu Leu Thr Ala  
 35 40 45  
 Ala Gln Leu Lys Ala Lys Tyr Leu Glu Asp Val Leu Glu Glu Leu  
 50 55 60  
 Thr Tyr Gly Pro Ala Pro Asp Leu Val Ile Ile Asn Ser Cys Leu  
 65 70 75  
 Trp Asp Leu Ser Arg Tyr Gly Arg Cys Ser Met Glu Ser Tyr Arg  
 80 85 90  
 Glu Asn Leu Glu Arg Val Phe Val Arg Met Asp Gln Val Leu Pro  
 95 100 105

Asp Ser Cys Leu	Leu Val Trp Asn Met	Ala Met Pro Leu Gly	Glu
110	115	120	
Arg Ile Thr Gly	Gly Phe Leu Leu Pro	Glu Leu Gln Pro Leu	Ala
125	130	135	
Gly Ser Leu Arg	Arg Asp Val Val Glu	Gly Asn Phe Tyr Ser	Ala
140	145	150	
Thr Leu Ala Gly	Asp His Cys Phe Asp	Val Leu Asp Leu His	Phe
155	160	165	
His Phe Arg His	Ala Val Gln His Arg	His Arg Asp Gly Val	His
170	175	180	
Trp Asp Gln His	Ala His Arg His Leu	Ser His Leu Leu Leu	Thr
185	190	195	
His Val Ala Asp	Ala Trp Gly Val Glu	Leu Pro Lys Arg Gly	Tyr
200	205	210	
Pro Pro Asp Pro	Trp Ile Glu Asp Trp	Ala Glu Met Asn His	Pro
215	220	225	
Phe Gln Gly Ser	His Arg Gln Thr Pro	Asp Phe Gly Glu His	Leu
230	235	240	
Ala Leu Leu Pro	Pro Pro Ser Ser	Leu Pro Pro Pro Met	Pro
245	250	255	
Phe Pro Tyr Pro	Leu Pro Gln Pro Ser	Pro Pro Pro Leu Phe	Pro
260	265	270	
Pro Leu Pro Gln	Asp Thr Pro Phe Phe	Pro Gly Gln Pro Phe	Pro
275	280	285	
Pro His Glu Phe	Phe Asn Tyr Asn Pro	Val Glu Asp Phe Ser	Met
290	295	300	
Pro Pro His Leu	Gly Cys Gly Pro Gly	Val Asn Phe Val Pro	Gly
305	310	315	
Pro Leu Pro Pro	Pro Ile Pro Gly Pro	Asn Pro His Gly Gln	His
320	325	330	
Trp Gly Pro Val	Val His Arg Gly Met	Pro Arg Tyr Val Pro	Asn
335	340	345	
Ser Pro Tyr His	Val Arg Arg Met Gly	Gly Pro Cys Arg Gln	Arg
350	355	360	
Leu Arg His Ser	Glu Arg Leu Ile His	Thr Tyr Lys Leu Asp	Arg
365	370	375	
Arg Pro Pro Ala	His Ser Gly Thr Trp	Pro Gly	
380	385		

&lt;210&gt; 4

&lt;211&gt; 833

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1674661CD1

&lt;400&gt; 4

Met Ala Pro His	Trp Ala Val Trp Leu Leu	Ala Ala Arg Leu Trp
1	5	10
Gly Leu Gly Ile	Gly Ala Glu Val Trp	Trp Asn Leu Val Pro Arg
20	25	30
Lys Thr Val Ser	Ser Gly Glu Leu Ala Thr	Val Val Arg Arg Phe
35	40	45
Ser Gln Thr Gly	Ile Gln Asp Phe Leu Thr	Leu Thr Leu Thr Glu
50	55	60
Pro Thr Gly Leu	Leu Tyr Val Gly Ala Arg	Glu Ala Leu Phe Ala
65	70	75
Phe Ser Met Glu	Ala Leu Glu Leu Gln Gly	Ala Ile Ser Trp Glu
80	85	90
Ala Pro Val Glu	Lys Lys Thr Glu Cys Ile	Gln Lys Gly Lys Asn
95	100	105
Asn Gln Thr Glu	Cys Phe Asn Phe Ile Arg	Phe Leu Gln Pro Tyr
110	115	120
Asn Ala Ser His	Leu Tyr Val Cys Gly Thr	Tyr Ala Phe Gln Pro
125	130	135

Lys	Cys	Thr	Tyr	Val	Asn	Met	Leu	Thr	Phe	Thr	Leu	Glu	His	Gly
				140					145					150
Glu	Phe	Glu	Asp	Gly	Lys	Gly	Lys	Cys	Pro	Tyr	Asp	Pro	Ala	Lys
				155					160					165
Gly	His	Ala	Gly	Leu	Leu	Val	Asp	Gly	Glu	Leu	Tyr	Ser	Ala	Thr
				170					175					180
Leu	Asn	Asn	Phe	Leu	Gly	Thr	Glu	Pro	Ile	Ile	Leu	Arg	Asn	Met
				185					190					195
Gly	Pro	His	His	Ser	Met	Lys	Thr	Glu	Tyr	Leu	Ala	Phe	Trp	Leu
				200					205					210
Asn	Glu	Pro	His	Phe	Val	Gly	Ser	Ala	Tyr	Val	Pro	Glu	Thr	Val
				215					220					225
Gly	Ser	Phe	Thr	Gly	Asp	Asp	Asp	Lys	Val	Tyr	Phe	Phe	Phe	Arg
				230					235					240
Glu	Arg	Ala	Leu	Glu	Ser	Asp	Cys	Tyr	Ala	Glu	Gln	Val	Val	Ala
				245					250					255
Arg	Val	Ala	Arg	Val	Cys	Lys	Gly	Asp	Met	Gly	Gly	Ala	Arg	Thr
				260					265					270
Leu	Gln	Arg	Lys	Trp	Thr	Thr	Phe	Leu	Lys	Ala	Arg	Leu	Ala	Cys
				275					280					285
Ser	Ala	Pro	Asn	Trp	Gln	Leu	Tyr	Phe	Asn	Gln	Leu	Gln	Ala	Met
				290					295					300
His	Thr	Leu	Gln	Asp	Thr	Ser	Trp	His	Asn	Thr	Thr	Phe	Phe	Gly
				305					310					315
Val	Phe	Gln	Ala	Gln	Trp	Gly	Asp	Met	Tyr	Leu	Ser	Ala	Ile	Cys
				320					325					330
Glu	Tyr	Gln	Leu	Glu	Glu	Ile	Gln	Arg	Val	Phe	Glu	Gly	Pro	Tyr
				335					340					345
Lys	Glu	Tyr	His	Glu	Glu	Ala	Gln	Lys	Trp	Asp	Arg	Tyr	Thr	Asp
				350					355					360
Pro	Val	Pro	Ser	Pro	Arg	Pro	Gly	Ser	Cys	Ile	Asn	Asn	Trp	His
				365					370					375
Arg	Arg	His	Gly	Tyr	Thr	Ser	Ser	Leu	Glu	Leu	Pro	Asp	Asn	Ile
				380					385					390
Leu	Asn	Phe	Val	Lys	Lys	His	Pro	Leu	Met	Glu	Glu	Gln	Val	Gly
				395					400					405
Pro	Arg	Trp	Ser	Arg	Pro	Leu	Leu	Val	Lys	Lys	Gly	Thr	Asn	Phe
				410					415					420
Thr	His	Leu	Val	Ala	Asp	Arg	Val	Thr	Gly	Leu	Asp	Gly	Ala	Thr
				425					430					435
Tyr	Thr	Val	Leu	Phe	Ile	Gly	Thr	Gly	Asp	Gly	Trp	Leu	Leu	Lys
				440					445					450
Ala	Val	Ser	Leu	Gly	Pro	Trp	Val	His	Leu	Ile	Glu	Glu	Leu	Gln
				455					460					465
Leu	Phe	Asp	Gln	Glu	Pro	Met	Arg	Ser	Leu	Val	Leu	Ser	Gln	Ser
				470					475					480
Lys	Lys	Leu	Leu	Phe	Ala	Gly	Ser	Arg	Ser	Gln	Leu	Val	Gln	Leu
				485					490					495
Pro	Val	Ala	Asp	Cys	Met	Lys	Tyr	Arg	Ser	Cys	Ala	Asp	Cys	Val
				500					505					510
Leu	Ala	Arg	Asp	Pro	Tyr	Cys	Ala	Trp	Ser	Val	Asn	Thr	Ser	Arg
				515					520					525
Cys	Val	Ala	Val	Gly	Gly	His	Ser	Gly	Ser	Leu	Leu	Ile	Gln	His
				530					535					540
Val	Met	Thr	Ser	Asp	Thr	Ser	Gly	Ile	Cys	Asn	Leu	Arg	Gly	Ser
				545					550					555
Lys	Lys	Val	Arg	Pro	Thr	Pro	Lys	Asn	Ile	Thr	Val	Val	Ala	Gly
				560					565					570
Thr	Asp	Leu	Val	Leu	Pro	Cys	His	Leu	Ser	Ser	Asn	Leu	Ala	His
				575					580					585
Ala	Arg	Trp	Thr	Phe	Gly	Gly	Arg	Asp	Leu	Pro	Ala	Glu	Gln	Pro
				590					595					600
Gly	Ser	Phe	Leu	Tyr	Asp	Ala	Arg	Leu	Gln	Ala	Leu	Val	Val	Met
				605					610					615
Ala	Ala	Gln	Pro	Arg	His	Ala	Gly	Ala	Tyr	His	Cys	Phe	Ser	Glu
				620					625					630
Glu	Gln	Gly	Ala	Arg	Leu	Ala	Ala	Glu	Gly	Tyr	Leu	Val	Ala	Val

Val	Ala	Gly	Pro	Ser	Val	Thr	Leu	Glu	Ala	Arg	Ala	Pro	Leu	Glu	635	640	645
Asn	Leu	Gly	Leu	Val	Trp	Leu	Ala	Val	Val	Ala	Leu	Gly	Ala	Val	650	655	660
Cys	Leu	Val	Leu	Leu	Leu	Leu	Val	Leu	Ser	Leu	Arg	Arg	Arg	Leu	665	670	675
Arg	Glu	Glu	Leu	Glu	Lys	Gly	Ala	Lys	Ala	Thr	Glu	Arg	Thr	Leu	680	685	690
Val	Tyr	Pro	Leu	Glu	Leu	Pro	Lys	Glu	Pro	Thr	Ser	Pro	Pro	Phe	695	700	705
Arg	Pro	Cys	Pro	Glu	Pro	Asp	Glu	Lys	Leu	Trp	Asp	Pro	Val	Gly	710	715	720
Tyr	Tyr	Tyr	Ser	Asp	Gly	Ser	Leu	Lys	Ile	Val	Pro	Gly	His	Ala	725	730	735
Arg	Cys	Gln	Pro	Gly	Gly	Gly	Pro	Pro	Ser	Pro	Pro	Pro	Gly	Ile	740	745	750
Pro	Gly	Gln	Pro	Leu	Pro	Ser	Pro	Thr	Arg	Leu	His	Leu	Gly	Gly	755	760	765
Gly	Arg	Asn	Ser	Asn	Ala	Asn	Gly	Tyr	Val	Arg	Leu	Gln	Leu	Gly	770	775	780
Gly	Glu	Asp	Arg	Gly	Gly	Leu	Gly	His	Pro	Leu	Pro	Glu	Leu	Ala	785	790	795
Asp	Glu	Leu	Arg	Arg	Lys	Leu	Gln	Gln	Arg	Gln	Pro	Leu	Pro	Asp	800	805	810
Ser	Asn	Pro	Glu	Glu	Ser	Ser	Val								815	820	825
															830		

<210> 5  
 <211> 410  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1689337CD1

Met	Ser	Asn	Ile	Ser	Leu	Leu	Ala	Leu	Phe	Ser	Ser	Gln	Val	Ser	<400> 5
1				5					10					15	
Ala	Ser	Leu	Lys	Ala	Leu	Ser	His	Phe	Phe	Ser	Leu	Cys	Phe	Arg	
				20					25					30	
Leu	Ala	Arg	Glu	Gln	Ala	Arg	Val	Cys	Glu	Leu	Gln	Ser	Gly	Asn	
				35					40					45	
Gln	Gln	Leu	Glu	Glu	Gln	Arg	Val	Glu	Leu	Val	Glu	Arg	Leu	Gln	
				50					55					60	
Ala	Met	Leu	Gln	Ala	His	Trp	Asp	Glu	Ala	Asn	Gln	Leu	Leu	Ser	
				65					70					75	
Thr	Thr	Leu	Pro	Pro	Pro	Asn	Pro	Pro	Ala	Pro	Pro	Ala	Gly	Pro	
				80					85					90	
Ser	Ser	Pro	Gly	Pro	Gln	Glu	Pro	Glu	Lys	Glu	Glu	Arg	Arg	Val	
				95					100					105	
Trp	Thr	Met	Pro	Pro	Met	Ala	Val	Ala	Leu	Lys	Pro	Val	Leu	Gln	
				110					115					120	
Gln	Ser	Arg	Glu	Ala	Arg	Asp	Glu	Leu	Pro	Gly	Ala	Pro	Pro	Val	
				125					130					135	
Leu	Cys	Ser	Ser	Ser	Ser	Asp	Leu	Ser	Leu	Leu	Gly	Pro	Ser		
				140					145					150	
Phe	Gln	Ser	Gln	His	Ser	Phe	Gln	Pro	Leu	Glu	Pro	Lys	Pro	Asp	
				155					160					165	
Leu	Thr	Ser	Ser	Thr	Ala	Gly	Ala	Phe	Ser	Ala	Leu	Gly	Ala	Phe	
				170					175					180	
His	Pro	Asp	His	Arg	Ala	Glu	Arg	Pro	Phe	Pro	Glu	Glu	Asp	Pro	
				185					190					195	
Gly	Pro	Asp	Gly	Glu	Gly	Leu	Leu	Lys	Gln	Gly	Leu	Pro	Pro	Ala	
				200					205					210	
Gln	Leu	Glu	Gly	Leu	Lys	Asn	Phe	Leu	His	Gln	Leu	Leu	Glu	Thr	

Val Pro Gln Asn	215	Val Asp Leu Leu Pro	225
Asn Glu Asn Pro Ser	230	Trp Glu Glu Ala Pro	240
Lys Ser Gly Pro	245	Lys Thr Lys Val Pro	255
Val Pro Arg Ile	260	Pro Glu Pro Pro Ser	270
Ala Met Ala Ser	275	Gly Ser Pro Glu Arg	285
His Ser Gln Gly	290	Leu Met Glu Val Ser	300
Gly Asp Gly Leu	305	Trp Gly Ala Leu Pro	315
Leu Leu Arg Leu	320	Leu Glu His Ser Gly	330
Glu Asp Leu Leu	335	Arg Arg Asn Thr Asp	345
Thr Asp Gly Arg	350	Ile Pro Ser Gln Ala	360
Arg Leu Gly Glu	365	Thr Glu Lys Pro Pro	375
Pro Arg Arg Leu	380	Ser Ser Met Arg Ser	390
Arg Lys Lys Ser	395	Gly Gly Val Trp Arg	405
	410		

<210> 6  
 <211> 360  
 <212> PRT  
 <213> Homo sapiens  
  
 <220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1746392CD1

<400> 6  
 Met Asp Thr Pro Leu Arg Arg Ser Arg Arg Leu Gly Gly Leu Arg  
 1 5 10 15  
 Pro Glu Ser Pro Glu Ser Leu Thr Ser Val Ser Arg Thr Arg Arg  
 20 25 30  
 Ala Leu Val Glu Phe Glu Ser Asn Pro Glu Glu Thr Arg Glu Pro  
 35 40 45  
 Gly Ser Pro Pro Ser Val Gln Arg Ala Gly Leu Gly Ser Pro Glu  
 50 55 60  
 Arg Pro Pro Lys Thr Ser Pro Gly Ser Pro Arg Leu Gln Gln Gly  
 65 70 75  
 Ala Gly Leu Glu Ser Pro Gln Gly Gln Pro Glu Pro Gly Ala Ala  
 80 85 90  
 Ser Pro Gln Arg Gln Gln Asp Leu His Leu Glu Ser Pro Gln Arg  
 95 100 105  
 Gln Pro Glu Tyr Ser Pro Glu Ser Pro Arg Cys Gln Pro Lys Pro  
 110 115 120  
 Ser Glu Glu Ala Pro Lys Cys Ser Gln Asp Gln Gly Val Leu Ala  
 125 130 135  
 Ser Glu Leu Ala Gln Asn Lys Glu Glu Leu Thr Pro Gly Ala Pro  
 140 145 150  
 Gln His Gln Leu Pro Pro Val Pro Gly Ser Pro Glu Pro Tyr Pro  
 155 160 165  
 Gly Gln Gln Ala Pro Gly Pro Glu Pro Ser Gln Pro Leu Leu Glu  
 170 175 180  
 Leu Thr Pro Arg Ala Pro Gly Ser Pro Arg Gly Gln His Glu Pro  
 185 190 195  
 Ser Lys Pro Pro Pro Ala Gly Glu Thr Val Thr Gly Gly Phe Gly  
 200 205 210  
 Ala Lys Lys Arg Lys Gly Ser Ser Ser Gln Ala Pro Ala Ser Lys

	215		220		225
Lys Leu Asn Lys	Glu Glu Leu Pro Val	Ile Pro Lys Gly Lys	Pro		
	230		235		240
Lys Ser Gly Arg	Val Trp Lys Asp Arg	Ser Lys Lys Arg Phe	Ser		
	245		250		255
Gln Met Leu Gln	Asp Lys Pro Leu Arg	Thr Ser Trp Gln Arg	Lys		
	260		265		270
Met Lys Glu Arg	Gln Glu Arg Lys Leu	Ala Lys Asp Phe Ala	Arg		
	275		280		285
His Leu Glu Glu	Glu Lys Glu Arg Arg	Arg Gln Glu Lys Lys	Gln		
	290		295		300
Arg Arg Ala Glu	Asn Leu Lys Arg Arg	Leu Glu Asn Glu Arg	Lys		
	305		310		315
Ala Glu Val Val	Gln Val Ile Arg Asn	Pro Ala Lys Leu Lys	Arg		
	320		325		330
Ala Lys Lys Lys	Gln Leu Arg Ser Ile	Glu Lys Arg Asp Thr	Leu		
	335		340		345
Ala Leu Leu Gln	Lys Gln Pro Pro Gln	Gln Pro Ala Ala Lys	Ile		
	350		355		360

&lt;210&gt; 7

&lt;211&gt; 377

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1825182CD1

&lt;400&gt; 7

Met Lys Thr Leu Pro	Leu Phe Val Cys	Ile Cys Ala Leu Ser	Ala
1	5	10	15
Cys Phe Ser Phe Ser	Glu Gly Arg Glu	Arg Asp His Glu Leu	Arg
	20	25	30
His Arg Arg His His	His Gln Ser Pro	Lys Ser His Phe Glu	Leu
	35	40	45
Pro His Tyr Pro Gly	Leu Leu Ala His	Gln Lys Pro Phe Ile	Arg
	50	55	60
Lys Ser Tyr Lys Cys	Leu His Lys Arg	Cys Arg Pro Lys Leu	Pro
	65	70	75
Pro Ser Pro Asn Lys	Pro Pro Lys Phe	Pro Asn Pro His Gln	Pro
	80	85	90
Pro Lys His Pro Asp	Lys Asn Ser Ser	Val Val Asn Pro Thr	Leu
	95	100	105
Val Ala Thr Thr Gln	Ile Pro Ser Val	Thr Phe Pro Ser Ala	Ser
	110	115	120
Thr Lys Ile Thr Thr	Leu Pro Asn Val	Thr Phe Leu Pro Gln	Asn
	125	130	135
Ala Thr Thr Ile Ser	Ser Arg Glu Asn	Val Asn Thr Ser Ser	Ser
	140	145	150
Val Ala Thr Leu Ala	Pro Val Asn Ser	Pro Ala Pro Gln Asp	Thr
	155	160	165
Thr Ala Ala Pro Pro	Thr Pro Ser Ala	Thr Thr Pro Ala Pro	Pro
	170	175	180
Ser Ser Ser Ala Pro	Pro Glu Thr Thr	Ala Ala Pro Pro Thr	Pro
	185	190	195
Ser Ala Thr Thr Gln	Ala Pro Pro Ser	Ser Ser Ala Pro Pro	Glu
	200	205	210
Thr Thr Ala Ala Pro	Pro Thr Pro Pro	Ala Thr Thr Pro Ala	Pro
	215	220	225
Pro Ser Ser Ser Ala	Pro Pro Glu Thr	Thr Ala Ala Pro Pro	Thr
	230	235	240
Pro Ser Ala Thr Thr	Pro Ala Pro Leu	Ser Ser Ser Ala Pro	Pro
	245	250	255
Glu Thr Thr Ala Val	Pro Pro Thr Pro	Ser Ala Thr Thr Leu	Asp
	260	265	270



```

Pro Ser Ser Ala Ser Ala Pro Pro Glu Thr Thr Ala Ala Pro Pro
      275      280      285
Thr Pro Ser Ala Thr Thr Pro Ala Pro Pro Ser Ser Pro Ala Pro
      290      295      300
Gln Glu Thr Thr Ala Ala Pro Ile Thr Thr Pro Asn Ser Ser Pro
      305      310      315
Thr Thr Leu Ala Pro Asp Thr Ser Glu Thr Ser Ala Ala Pro Thr
      320      325      330
His Gln Thr Thr Thr Ser Val Thr Thr Gln Thr Thr Thr Thr Lys
      335      340      345
Gln Pro Thr Ser Ala Pro Gly Gln Asn Lys Ile Ser Arg Phe Leu
      350      355      360
Leu Tyr Met Lys Asn Leu Leu Asn Arg Ile Ile Asp Asp Met Val
      365      370      375
Glu Gln

```

<210> 8  
 <211> 182  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2155541CD1

```

<400> 8
Met Leu Val Leu Val Leu Gly Asp Leu His Ile Pro His Arg Cys
  1      5      10      15
Asn Ser Leu Pro Ala Lys Phe Lys Lys Leu Leu Val Pro Gly Lys
      20      25      30
Ile Gln His Ile Leu Cys Thr Gly Asn Leu Cys Thr Lys Glu Ser
      35      40      45
Tyr Asp Tyr Leu Lys Thr Leu Ala Gly Asp Val His Ile Val Arg
      50      55      60
Gly Asp Phe Asp Glu Asn Leu Asn Tyr Pro Glu Gln Lys Val Val
      65      70      75
Thr Val Gly Gln Phe Lys Ile Gly Leu Ile His Gly His Gln Val
      80      85      90
Ile Pro Trp Gly Asp Met Ala Ser Leu Ala Leu Leu Gln Arg Gln
      95      100      105
Phe Asp Val Asp Ile Leu Ile Ser Gly His Thr His Lys Phe Glu
      110      115      120
Ala Phe Glu His Glu Asn Lys Phe Tyr Ile Asn Pro Gly Ser Ala
      125      130      135
Thr Gly Ala Tyr Asn Ala Leu Glu Thr Asn Ile Ile Pro Ser Phe
      140      145      150
Val Leu Met Asp Ile Gln Ala Ser Thr Val Val Thr Tyr Val Tyr
      155      160      165
Gln Leu Ile Gly Asp Asp Val Lys Val Glu Arg Ile Glu Tyr Lys
      170      175      180
Lys Pro

```

<210> 9  
 <211> 513  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2215706CD1

```

<400> 9
Met Asn Met Asn Phe Gly Asp Trp His Leu Phe Arg Ser Thr Val
  1      5      10      15
Leu Glu Met Arg Asn Ala Glu Ser His Val Val Pro Glu Asp Pro

```

9/42

<210> 10  
 <211> 361  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2347692CD1

<400> 10  
 Met Tyr Gly Lys Gly Lys Ser Asn Ser Ser Ala Val Pro Ser Asp  
 1 5 10 15  
 Ser Gln Ala Arg Glu Lys Leu Ala Leu Tyr Val Tyr Glu Tyr Leu  
 20 25 30  
 Leu His Val Gly Ala Gln Lys Ser Ala Gln Thr Phe Leu Ser Glu  
 35 40 45  
 Ile Arg Trp Glu Lys Asn Ile Thr Leu Gly Glu Pro Pro Gly Phe  
 50 55 60  
 Leu His Ser Trp Trp Cys Val Phe Trp Asp Leu Tyr Cys Ala Ala  
 65 70 75  
 Pro Glu Arg Arg Glu Thr Cys Glu His Ser Ser Glu Ala Lys Ala  
 80 85 90  
 Phe His Asp Tyr Ser Ala Ala Ala Ala Pro Ser Pro Val Leu Gly  
 95 100 105  
 Asn Ile Pro Pro Gly Asp Gly Met Pro Val Gly Pro Val Pro Pro  
 110 115 120  
 Gly Phe Phe Gln Pro Phe Met Ser Pro Arg Tyr Pro Gly Gly Pro  
 125 130 135  
 Arg Pro Pro Leu Arg Ile Pro Asn Gln Ala Leu Gly Gly Val Pro  
 140 145 150  
 Gly Ser Gln Pro Leu Leu Pro Ser Gly Met Asp Pro Thr Arg Gln  
 155 160 165  
 Gln Gly His Pro Asn Met Gly Gly Pro Met Gln Arg Met Thr Pro  
 170 175 180  
 Pro Arg Gly Met Val Pro Leu Gly Pro Gln Asn Tyr Gly Gly Ala  
 185 190 195  
 Met Arg Pro Pro Leu Asn Ala Leu Gly Gly Pro Gly Met Pro Gly  
 200 205 210  
 Met Asn Met Gly Pro Gly Gly Gly Arg Pro Trp Pro Asn Pro Thr  
 215 220 225  
 Asn Ala Asn Ser Ile Pro Tyr Ser Ser Ala Ser Pro Gly Asn Tyr  
 230 235 240  
 Val Gly Pro Pro Gly Gly Gly Gly Pro Pro Gly Thr Pro Ile Met  
 245 250 255  
 Pro Ser Pro Ala Asp Ser Thr Asn Ser Gly Asp Asn Met Tyr Thr  
 260 265 270  
 Leu Met Asn Ala Val Pro Pro Gly Pro Asn Arg Pro Asn Phe Pro  
 275 280 285  
 Met Gly Pro Gly Ser Asp Gly Pro Met Gly Gly Leu Gly Gly Met  
 290 295 300  
 Glu Ser His His Met Asn Gly Ser Leu Gly Ser Gly Asp Met Asp  
 305 310 315  
 Ser Ile Ser Lys Asn Ser Pro Asn Asn Met Ser Leu Ser Asn Gln  
 320 325 330  
 Pro Gly Thr Pro Arg Asp Asp Gly Glu Met Gly Gly Asn Phe Leu  
 335 340 345  
 Asn Pro Phe Gln Ser Glu Ser Tyr Ser Pro Ser Met Thr Met Ser  
 350 355 360  
 Val

<210> 11  
 <211> 327  
 <212> PRT  
 <213> Homo sapiens

<220>

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2579048CD1

&lt;400&gt; 11

Met	Ala	Leu	Val	His	Lys	Leu	Leu	His	Gly	Thr	Tyr	Phe	Leu	Arg
1				5					10					15
Lys	Phe	Ser	Lys	Pro	Thr	Ser	Ala	Leu	Tyr	Pro	Phe	Leu	Gly	Ile
			20						25					30
Leu	Phe	Ala	Glu	Tyr	Ser	Ser	Ser	Leu	Gln	Lys	Pro	Val	Ala	Ser
			35						40					45
Pro	Gly	Lys	Ala	Ser	Ser	Gln	Arg	Lys	Thr	Glu	Gly	Asp	Leu	Gln
			50						55					60
Gly	Asp	His	Gln	Lys	Glu	Val	Ala	Leu	Asp	Ile	Thr	Ser	Ser	Glu
			65						70					75
Glu	Lys	Pro	Asp	Val	Ser	Phe	Asp	Lys	Ala	Ile	Arg	Asp	Glu	Ala
			80						85					90
Ile	Tyr	His	Phe	Arg	Leu	Leu	Lys	Asp	Glu	Ile	Val	Asp	His	Trp
			95						100					105
Arg	Gly	Pro	Glu	Gly	His	Pro	Leu	His	Glu	Val	Leu	Leu	Glu	Gln
			110						115					120
Ala	Lys	Val	Val	Trp	Gln	Phe	Arg	Gly	Lys	Glu	Asp	Leu	Asp	Lys
			125						130					135
Trp	Thr	Val	Thr	Ser	Asp	Lys	Thr	Ile	Gly	Gly	Arg	Ser	Glu	Val
			140						145					150
Phe	Leu	Lys	Met	Gly	Lys	Asn	Asn	Gln	Ser	Ala	Leu	Leu	Tyr	Gly
			155						160					165
Thr	Leu	Ser	Ser	Glu	Ala	Pro	Gln	Asp	Gly	Glu	Ser	Thr	Arg	Ser
			170						175					180
Gly	Tyr	Cys	Ala	Met	Ile	Ser	Arg	Ile	Pro	Arg	Gly	Ala	Phe	Glu
			185						190					195
Arg	Lys	Met	Ser	Tyr	Asp	Trp	Ser	Gln	Phe	Asn	Thr	Leu	Tyr	Leu
			200						205					210
Arg	Val	Arg	Gly	Asp	Gly	Arg	Pro	Trp	Met	Val	Asn	Ile	Lys	Glu
			215						220					225
Asp	Thr	Asp	Phe	Phe	Gln	Arg	Thr	Asn	Gln	Met	Tyr	Ser	Tyr	Phe
			230						235					240
Met	Phe	Thr	Arg	Gly	Gly	Pro	Tyr	Trp	Gln	Glu	Val	Lys	Ile	Pro
			245						250					255
Phe	Ser	Lys	Phe	Phe	Phe	Ser	Asn	Arg	Gly	Arg	Ile	Arg	Asp	Val
			260						265					270
Gln	His	Glu	Leu	Pro	Leu	Asp	Lys	Ile	Ser	Ser	Ile	Gly	Phe	Thr
			275						280					285
Leu	Ala	Asp	Lys	Val	Asp	Gly	Pro	Phe	Phe	Leu	Glu	Ile	Asp	Phe
			290						295					300
Ile	Gly	Val	Phe	Thr	Asp	Pro	Ala	His	Thr	Glu	Glu	Phe	Ala	Tyr
			305						310					315
Glu	Asn	Ser	Pro	Glu	Leu	Asn	Pro	Arg	Leu	Phe	Lys			
			320						325					

&lt;210&gt; 12

&lt;211&gt; 1110

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2604493CD1

&lt;400&gt; 12

Met	Pro	Ala	Pro	Glu	Gln	Ala	Ser	Leu	Val	Glu	Glu	Gly	Gln	Pro
1				5					10					15
Gln	Thr	Arg	Gln	Glu	Ala	Ala	Ser	Thr	Gly	Pro	Gly	Met	Glu	Pro
			20						25					30
Glu	Thr	Thr	Ala	Thr	Thr	Ile	Leu	Ala	Ser	Val	Lys	Glu	Gln	Glu
			35						40					45
Leu	Gln	Phe	Gln	Arg	Leu	Thr	Arg	Glu	Leu	Glu	Val	Glu	Arg	Gln
			50						55					60

Ile	Val	Ala	Ser	Gln	Leu	Glu	Arg	Cys	Arg	Leu	Gly	Ala	Glu	Ser
				65					70					75
Pro	Ser	Ile	Ala	Ser	Thr	Ser	Ser	Thr	Glu	Lys	Ser	Phe	Pro	Trp
				80					85					90
Arg	Ser	Thr	Asp	Val	Pro	Asn	Thr	Gly	Val	Ser	Lys	Pro	Arg	Val
				95					100					105
Ser	Asp	Ala	Val	Gln	Pro	Asn	Asn	Tyr	Leu	Ile	Arg	Thr	Glu	Pro
				110					115					120
Glu	Gln	Gly	Thr	Leu	Tyr	Ser	Pro	Glu	Gln	Thr	Ser	Leu	His	Glu
				125					130					135
Ser	Glu	Gly	Ser	Leu	Gly	Asn	Ser	Arg	Ser	Ser	Thr	Gln	Met	Asn
				140					145					150
Ser	Tyr	Ser	Asp	Ser	Gly	Tyr	Gln	Glu	Ala	Gly	Ser	Phe	His	Asn
				155					160					165
Ser	Gln	Asn	Val	Ser	Lys	Ala	Asp	Asn	Arg	Gln	Gln	His	Ser	Phe
				170					175					180
Ile	Gly	Ser	Thr	Asn	Asn	His	Val	Val	Arg	Asn	Ser	Arg	Ala	Glu
				185					190					195
Gly	Gln	Thr	Leu	Val	Gln	Pro	Ser	Val	Ala	Asn	Arg	Ala	Met	Arg
				200					205					210
Arg	Val	Ser	Ser	Val	Pro	Ser	Arg	Ala	Gln	Ser	Pro	Ser	Tyr	Val
				215					220					225
Ile	Ser	Thr	Gly	Val	Ser	Pro	Ser	Arg	Gly	Ser	Leu	Arg	Thr	Ser
				230					235					240
Leu	Gly	Ser	Gly	Phe	Gly	Ser	Pro	Ser	Val	Thr	Asp	Pro	Arg	Pro
				245					250					255
Leu	Asn	Pro	Ser	Ala	Tyr	Ser	Ser	Thr	Thr	Leu	Pro	Ala	Ala	Arg
				260					265					270
Ala	Ala	Ser	Pro	Tyr	Ser	Gln	Arg	Pro	Ala	Ser	Pro	Thr	Ala	Ile
				275					280					285
Arg	Arg	Ile	Gly	Ser	Val	Thr	Ser	Arg	Gln	Thr	Ser	Asn	Pro	Asn
				290					295					300
Gly	Pro	Thr	Pro	Gln	Tyr	Gln	Thr	Thr	Ala	Arg	Val	Gly	Ser	Pro
				305					310					315
Leu	Thr	Leu	Thr	Asp	Ala	Gln	Thr	Arg	Val	Ala	Ser	Pro	Ser	Gln
				320					325					330
Gly	Gln	Val	Gly	Ser	Ser	Ser	Pro	Lys	Arg	Ser	Gly	Met	Thr	Ala
				335					340					345
Val	Pro	Gln	His	Leu	Gly	Pro	Ser	Leu	Gln	Arg	Thr	Val	His	Asp
				350					355					360
Met	Glu	Gln	Phe	Gly	Gln	Gln	Gln	Tyr	Asp	Ile	Tyr	Glu	Arg	Met
				365					370					375
Val	Pro	Pro	Arg	Pro	Asp	Ser	Leu	Thr	Gly	Leu	Arg	Ser	Ser	Tyr
				380					385					390
Ala	Ser	Gln	His	Ser	Gln	Leu	Gly	Gln	Asp	Leu	Arg	Ser	Ala	Val
				395					400					405
Ser	Pro	Asp	Leu	His	Ile	Thr	Pro	Ile	Tyr	Glu	Gly	Arg	Thr	Tyr
				410					415					420
Tyr	Ser	Pro	Val	Tyr	Arg	Ser	Pro	Asn	His	Gly	Thr	Val	Glu	Leu
				425					430					435
Gln	Gly	Ser	Gln	Thr	Ala	Leu	Tyr	Arg	Thr	Gly	Ser	Gly	Ile	Gly
				440					445					450
Asn	Leu	Gln	Arg	Thr	Ser	Ser	Gln	Arg	Ser	Thr	Leu	Thr	Tyr	Gln
				455					460					465
Arg	Asn	Asn	Tyr	Ala	Leu	Asn	Thr	Thr	Ala	Thr	Tyr	Ala	Glu	Pro
				470					475					480
Tyr	Arg	Pro	Ile	Gln	Tyr	Arg	Val	Gln	Glu	Cys	Asn	Tyr	Asn	Arg
				485					490					495
Leu	Gln	His	Ala	Val	Pro	Ala	Asp	Asp	Gly	Thr	Thr	Arg	Ser	Pro
				500					505					510
Ser	Ile	Asp	Ser	Ile	Gln	Lys	Asp	Pro	Arg	Glu	Phe	Ala	Trp	Arg
				515					520					525
Asp	Pro	Glu	Leu	Pro	Glu	Val	Ile	His	Met	Leu	Gln	His	Gln	Phe
				530					535					540
Pro	Ser	Val	Gln	Ala	Asn	Ala	Ala	Ala	Tyr	Leu	Gln	His	Leu	Cys
				545					550					555
Phe	Gly	Asp	Asn	Lys	Val	Lys	Met	Glu	Val	Cys	Arg	Leu	Gly	Gly

Ile	Lys	His	Leu	Val	Asp	Leu	Leu	Asp	His	Arg	Val	Leu	Glu	Val
				560					565					570
Gln	Lys	Asn	Ala	Cys	Gly	Ala	Leu	Arg	Asn	Leu	Val	Phe	Gly	Lys
				575					580					585
Ser	Thr	Asp	Glu	Asn	Lys	Ile	Ala	Met	Lys	Asn	Val	Gly	Gly	Ile
				590					595					600
Pro	Ala	Leu	Leu	Arg	Leu	Leu	Arg	Lys	Ser	Ile	Asp	Ala	Glu	Val
				605					610					615
Arg	Glu	Leu	Val	Thr	Gly	Val	Leu	Trp	Asn	Leu	Ser	Ser	Cys	Asp
				620					625					630
Ala	Val	Lys	Met	Thr	Ile	Ile	Arg	Asp	Ala	Leu	Ser	Thr	Leu	Thr
				635					640					645
Asn	Thr	Val	Ile	Val	Pro	His	Ser	Gly	Trp	Asn	Asn	Ser	Ser	Phe
				650					655					660
Asp	Asp	Asp	His	Lys	Ile	Lys	Phe	Gln	Thr	Ser	Leu	Val	Leu	Arg
				665					670					675
Asn	Thr	Thr	Gly	Cys	Leu	Arg	Asn	Leu	Ser	Ser	Ala	Gly	Glu	Glu
				680					685					690
Ala	Arg	Lys	Gln	Met	Arg	Ser	Cys	Glu	Gly	Leu	Val	Asp	Ser	Leu
				695					700					705
Leu	Tyr	Val	Ile	His	Thr	Cys	Val	Asn	Thr	Ser	Asp	Tyr	Asp	Ser
				710					715					720
Lys	Thr	Val	Glu	Asn	Cys	Val	Cys	Thr	Leu	Arg	Asn	Leu	Ser	Tyr
				725					730					735
Arg	Leu	Glu	Leu	Glu	Val	Pro	Gln	Ala	Arg	Leu	Leu	Gly	Leu	Asn
				740					745					750
Glu	Leu	Asp	Asp	Leu	Leu	Gly	Lys	Glu	Ser	Pro	Ser	Lys	Asp	Ser
				755					760					765
Glu	Pro	Ser	Cys	Trp	Gly	Lys	Lys	Lys	Lys	Lys	Lys	Lys	Arg	Thr
				770					775					780
Pro	Gln	Glu	Asp	Gln	Trp	Asp	Gly	Val	Gly	Pro	Ile	Pro	Gly	Leu
				785					790					795
Ser	Lys	Ser	Pro	Lys	Gly	Val	Glu	Met	Leu	Trp	His	Pro	Ser	Val
				800					805					810
Val	Lys	Pro	Tyr	Leu	Thr	Leu	Leu	Ala	Glu	Ser	Ser	Asn	Pro	Ala
				815					820					825
Thr	Leu	Glu	Gly	Ser	Ala	Gly	Ser	Leu	Gln	Asn	Leu	Ser	Ala	Gly
				830					835					840
Asn	Trp	Lys	Phe	Ala	Ala	Tyr	Ile	Arg	Ala	Ala	Val	Arg	Lys	Glu
				845					850					855
Lys	Gly	Leu	Pro	Ile	Leu	Val	Glu	Leu	Leu	Arg	Met	Asp	Asn	Asp
				860					865					870
Arg	Val	Val	Ser	Ser	Val	Ala	Thr	Ala	Leu	Arg	Asn	Met	Ala	Leu
				875					880					885
Asp	Val	Arg	Asn	Lys	Glu	Leu	Ile	Gly	Lys	Tyr	Ala	Met	Arg	Asp
				890					895					900
Leu	Val	Asn	Arg	Leu	Pro	Gly	Gly	Asn	Gly	Pro	Ser	Val	Leu	Ser
				905					910					915
Asp	Glu	Thr	Met	Ala	Ala	Ile	Cys	Cys	Ala	Leu	His	Glu	Val	Thr
				920					925					930
Ser	Lys	Asn	Met	Glu	Asn	Ala	Lys	Ala	Leu	Ala	Asp	Ser	Gly	Gly
				935					940					945
Ile	Glu	Lys	Leu	Val	Asn	Ile	Thr	Lys	Gly	Arg	Gly	Asp	Arg	Ser
				950					955					960
Ser	Leu	Lys	Val	Val	Lys	Ala	Ala	Ala	Gln	Val	Leu	Asn	Thr	Leu
				965					970					975
Trp	Gln	Tyr	Arg	Asp	Leu	Arg	Ser	Ile	Tyr	Lys	Lys	Asp	Gly	Trp
				980					985					990
Asn	Gln	Asn	His	Phe	Ile	Thr	Pro	Val	Ser	Thr	Leu	Glu	Arg	Asp
				995					1000					1005
Arg	Phe	Lys	Ser	His	Pro	Ser	Leu	Ser	Thr	Thr	Asn	Gln	Gln	Met
				1010					1015					1020
Ser	Pro	Ile	Ile	Gln	Ser	Val	Gly	Ser	Thr	Ser	Ser	Ser	Pro	Ala
				1025					1030					1035
Leu	Leu	Gly	Ile	Arg	Asp	Pro	Arg	Ser	Glu	Tyr	Asp	Arg	Thr	Gln
				1040					1045					1050
				1055					1060					1065

Pro Pro Met Gln Tyr Tyr Asn Ser Gln Gly Asp Ala Thr His Lys  
 1070 1075 1080  
 Gly Leu Tyr Pro Gly Lys Thr Pro Val Gly Cys Val Ile Gln Ser  
 1085 1090 1095  
 Leu Glu Lys Pro His Phe Gln Ala Leu Gly Gln Trp Pro Gly Lys  
 1100 1105 1110

<210> 13  
 <211> 386  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2787182CD1

<400> 13  
 Met Asp Arg Phe Val Trp Thr Ser Gly Leu Leu Glu Ile Asn Glu  
 1 5 10 15  
 Thr Leu Val Ile Gln Gln Arg Gly Val Arg Ile Tyr Asp Gly Glu  
 20 25 30  
 Glu Lys Ile Lys Phe Asp Ala Gly Thr Leu Leu Leu Ser Thr His  
 35 40 45  
 Arg Leu Ile Trp Arg Asp Gln Lys Asn His Glu Cys Cys Met Ala  
 50 55 60  
 Ile Leu Leu Ser Gln Ile Val Phe Ile Glu Glu Gln Ala Ala Gly  
 65 70 75  
 Ile Gly Lys Ser Ala Lys Ile Val Val His Leu His Pro Ala Pro  
 80 85 90  
 Pro Asn Lys Glu Pro Gly Pro Phe Gln Ser Ser Lys Asn Ser Tyr  
 95 100 105  
 Ile Lys Leu Ser Phe Lys Glu His Gly Gln Ile Glu Phe Tyr Arg  
 110 115 120  
 Arg Leu Ser Glu Glu Met Thr Gln Arg Arg Trp Glu Asn Met Pro  
 125 130 135  
 Val Ser Gln Ser Leu Gln Thr Asn Arg Gly Pro Gln Pro Gly Arg  
 140 145 150  
 Ile Arg Ala Val Gly Ile Val Gly Ile Glu Arg Lys Leu Glu Glu  
 155 160 165  
 Lys Lys Lys Glu Thr Asp Lys Asn Ile Ser Glu Ala Phe Glu Asp  
 170 175 180  
 Leu Ser Lys Leu Met Ile Lys Ala Lys Glu Met Val Glu Leu Ser  
 185 190 195  
 Lys Ser Ile Ala Asn Lys Ile Lys Asp Lys Gln Gly Asp Ile Thr  
 200 205 210  
 Glu Asp Glu Thr Ile Arg Phe Lys Ser Tyr Leu Leu Ser Met Gly  
 215 220 225  
 Ile Ala Asn Pro Val Thr Arg Glu Thr Tyr Gly Ser Gly Thr Gln  
 230 235 240  
 Tyr His Met Gln Leu Ala Lys Gln Leu Ala Gly Ile Leu Gln Val  
 245 250 255  
 Pro Leu Glu Glu Arg Gly Gly Ile Met Ser Leu Thr Glu Val Tyr  
 260 265 270  
 Cys Leu Val Asn Arg Ala Arg Gly Met Glu Leu Leu Ser Pro Glu  
 275 280 285  
 Asp Leu Val Asn Ala Cys Lys Met Leu Glu Ala Leu Lys Leu Pro  
 290 295 300  
 Leu Arg Leu Arg Val Phe Asp Ser Gly Val Met Val Ile Glu Leu  
 305 310 315  
 Gln Ser His Lys Glu Glu Glu Met Val Ala Ser Ala Leu Glu Thr  
 320 325 330  
 Val Ser Glu Lys Gly Ser Leu Thr Ser Glu Glu Phe Ala Lys Leu  
 335 340 345  
 Val Gly Met Ser Val Leu Leu Ala Lys Glu Arg Leu Leu Leu Ala  
 350 355 360  
 Glu Lys Met Gly His Leu Cys Arg Asp Ser Val Glu Gly Leu

365  
Arg Phe Tyr Pro Asn Leu Phe Met Thr Gln Ser  
380 385

375

<210> 14  
<211> 181  
<212> PRT  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 3096668CD1

<400> 14  
Met Glu Val Val Glu Ala Ala Ala Ala Gln Leu Glu Thr Leu Lys  
1 5 10 15  
Phe Asn Gly Thr Asp Phe Gly Val Gly Glu Gly Pro Ala Ala Pro  
20 25 30  
Ser Pro Gly Ser Ala Pro Val Pro Gly Thr Gln Pro Pro Leu Gln  
35 40 45  
Ser Phe Glu Gly Ser Pro Asp Ala Gly Gln Thr Val Glu Val Lys  
50 55 60  
Pro Ala Gly Glu Gln Pro Leu Gln Pro Val Leu Asn Ala Val Ala  
65 70 75  
Ala Gly Thr Pro Ala Pro Gln Pro Gln Pro Pro Ala Glu Ser Pro  
80 85 90  
Ala Cys Gly Asp Cys Val Thr Ser Pro Gly Ala Ala Glu Pro Ala  
95 100 105  
Arg Ala Pro Asp Ser Leu Glu Thr Ser Asp Ser Asp Ser Asp Ser  
110 115 120  
Asp Ser Glu Thr Asp Ser Asp Ser Ser Ser Ser Ser Ser Ser  
125 130 135  
Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Cys Ile Ser Leu Pro  
140 145 150  
Pro Val Leu Ser Asp Gly Asp Asp Asp Leu Gln Val Glu Lys Glu  
155 160 165  
Asn Lys Asn Phe Pro Leu Lys Thr Lys Asp Glu Leu Leu Leu Asn  
170 175 180  
Leu

<210> 15  
<211> 374  
<212> PRT  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 3143411CD1

<400> 15  
Met Arg Pro Gly Thr Ala Leu Gln Ala Val Leu Leu Ala Val Leu  
1 5 10 15  
Leu Val Gly Leu Arg Ala Ala Thr Gly Arg Leu Leu Ser Gly Gln  
20 25 30  
Pro Val Cys Arg Gly Gly Thr Gln Arg Pro Cys Tyr Lys Val Ile  
35 40 45  
Tyr Phe His Asp Thr Ser Arg Arg Leu Asn Phe Glu Glu Ala Lys  
50 55 60  
Glu Ala Cys Arg Arg Asp Gly Gly Gln Leu Val Ser Ile Glu Ser  
65 70 75  
Glu Asp Glu Gln Lys Leu Ile Glu Lys Phe Ile Glu Asn Leu Leu  
80 85 90  
Pro Ser Asp Gly Asp Phe Trp Ile Gly Leu Arg Arg Arg Glu Glu  
95 100 105  
Lys Gln Ser Asn Ser Thr Ala Cys Gln Asp Leu Tyr Ala Trp Thr  
110 115 120



Asp	Gly	Ser	Ile	Ser	Gln	Phe	Arg	Asn	Trp	Tyr	Val	Asp	Glu	Pro	
				125					130					135	
Ser	Cys	Gly	Ser	Glu	Val	Cys	Val	Val	Met	Tyr	His	Gln	Pro	Ser	
				140					145					150	
Ala	Pro	Ala	Gly	Ile	Gly	Gly	Pro	Tyr	Met	Phe	Gln	Trp	Asn	Asp	
				155					160					165	
Asp	Arg	Cys	Asn	Met	Lys	Asn	Asn	Phe	Ile	Cys	Lys	Tyr	Ser	Asp	
				170					175					180	
Glu	Lys	Pro	Ala	Val	Pro	Ser	Arg	Glu	Ala	Glu	Gly	Glu	Glu	Thr	
				185					190					195	
Glu	Leu	Thr	Thr	Pro	Val	Leu	Pro	Glu	Glu	Thr	Gln	Glu	Glu	Asp	
				200					205					210	
Ala	Lys	Lys	Thr	Phe	Lys	Glu	Ser	Arg	Glu	Ala	Ala	Leu	Asn	Leu	
				215					220					225	
Ala	Tyr	Ile	Leu	Ile	Pro	Ser	Ile	Pro	Leu	Leu	Leu	Leu	Leu	Val	
				230					235					240	
Val	Thr	Thr	Val	Val	Cys	Trp	Val	Trp	Ile	Cys	Arg	Lys	Arg	Lys	
				245					250					255	
Arg	Glu	Gln	Pro	Asp	Pro	Ser	Thr	Lys	Lys	Gln	His	Thr	Ile	Trp	
				260					265					270	
Pro	Ser	Pro	His	Gln	Gly	Asn	Ser	Pro	Asp	Leu	Glu	Val	Tyr	Asn	
				275					280					285	
Val	Ile	Arg	Lys	Gln	Ser	Glu	Ala	Asp	Leu	Ala	Glu	Thr	Arg	Pro	
				290					295					300	
Asp	Leu	Lys	Asn	Ile	Ser	Phe	Arg	Val	Cys	Ser	Gly	Glu	Ala	Thr	
				305					310					315	
Pro	Asp	Asp	Met	Ser	Cys	Asp	Tyr	Asp	Asn	Met	Ala	Val	Asn	Pro	
				320					325					330	
Ser	Glu	Ser	Gly	Phe	Val	Thr	Leu	Val	Ser	Val	Glu	Ser	Gly	Phe	
				335					340					345	
Val	Thr	Asn	Asp	Ile	Tyr	Glu	Phe	Ser	Pro	Asp	Gln	Met	Gly	Arg	
				350					355					360	
Ser	Lys	Glu	Ser	Gly	Trp	Val	Glu	Asn	Glu	Ile	Tyr	Gly	Tyr		
				365					370						

<210> 16  
 <211> 102  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3170835CD1

<400> 16															
Met	Lys	Phe	Ala	Ile	Val	Leu	Phe	Ala	Leu	Phe	Ala	Val	Ala	Leu	
1				5					10					15	
Ala	Ala	Pro	Thr	Val	Glu	Val	Leu	Arg	Ser	Asp	Ser	Asn	Val	Gly	
				20					25					30	
Ile	Asp	Asn	Tyr	Ser	Tyr	Ala	Val	Glu	Thr	Ser	Asp	Gly	Thr	Ser	
				35					40					45	
Lys	Ser	Glu	Glu	Gly	Val	Leu	Lys	Asn	Ala	Gly	Thr	Glu	Leu	Glu	
				50					55					60	
Ala	Ile	Ser	Thr	His	Gly	Ser	Phe	Ser	Tyr	Val	Gly	Pro	Asp	Gly	
				65					70					75	
Gln	Thr	Tyr	Thr	Val	Thr	Tyr	Val	Ala	Asp	Glu	Asn	Gly	Phe	Gln	
				80					85					90	
Pro	Gln	Gly	Ala	His	Leu	Pro	Val	Ala	Pro	Val	Ala				
				95					100						

<210> 17  
 <211> 510  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature

&lt;223&gt; Incyte ID No: 3550808CD1

&lt;400&gt; 17

Met	Arg	Pro	Gly	Leu	Ser	Phe	Leu	Leu	Ala	Leu	Leu	Phe	Phe	Leu
1				5					10					15
Gly	Gln	Ala	Ala	Gly	Asp	Leu	Gly	Asp	Val	Gly	Pro	Pro	Ile	Pro
				20					25					30
Ser	Pro	Gly	Phe	Ser	Ser	Phe	Pro	Gly	Val	Asp	Ser	Ser	Ser	Ser
				35					40					45
Phe	Ser	Ser	Ser	Ser	Arg	Ser	Gly	Ser	Ser	Ser	Arg	Ser	Ser	Leu
				50					55					60
Gly	Ser	Gly	Gly	Ser	Val	Ser	Gln	Leu	Phe	Ser	Asn	Phe	Thr	Gly
				65					70					75
Ser	Val	Asp	Asp	Arg	Gly	Thr	Cys	Gln	Cys	Ser	Val	Ser	Leu	Pro
				80					85					90
Asp	Thr	Thr	Phe	Pro	Val	Asp	Arg	Val	Glu	Arg	Leu	Glu	Phe	Thr
				95					100					105
Ala	His	Val	Leu	Ser	Gln	Lys	Phe	Glu	Lys	Glu	Leu	Ser	Lys	Val
				110					115					120
Arg	Glu	Tyr	Val	Gln	Leu	Ile	Ser	Val	Tyr	Glu	Lys	Lys	Leu	Leu
				125					130					135
Asn	Leu	Thr	Val	Arg	Ile	Asp	Ile	Met	Glu	Lys	Asp	Thr	Ile	Ser
				140					145					150
Tyr	Thr	Glu	Leu	Asp	Phe	Glu	Leu	Ile	Lys	Val	Glu	Val	Lys	Glu
				155					160					165
Met	Glu	Lys	Leu	Val	Ile	Gln	Leu	Lys	Glu	Ser	Phe	Gly	Gly	Ser
				170					175					180
Ser	Glu	Ile	Val	Asp	Gln	Leu	Glu	Val	Glu	Ile	Arg	Asn	Met	Thr
				185					190					195
Leu	Leu	Val	Glu	Lys	Leu	Glu	Thr	Leu	Asp	Lys	Asn	Asn	Val	Leu
				200					205					210
Ala	Ile	Arg	Arg	Glu	Ile	Val	Ala	Leu	Lys	Thr	Lys	Leu	Lys	Glu
				215					220					225
Cys	Glu	Ala	Ser	Lys	Asp	Gln	Asn	Thr	Pro	Val	Val	His	Pro	Pro
				230					235					240
Pro	Thr	Pro	Gly	Ser	Cys	Gly	His	Gly	Gly	Val	Val	Asn	Ile	Ser
				245					250					255
Lys	Pro	Ser	Val	Val	Gln	Leu	Asn	Trp	Arg	Gly	Phe	Ser	Tyr	Leu
				260					265					270
Tyr	Gly	Ala	Trp	Gly	Arg	Asp	Tyr	Ser	Pro	Gln	His	Pro	Asn	Lys
				275					280					285
Gly	Leu	Tyr	Trp	Val	Ala	Pro	Leu	Asn	Thr	Asp	Gly	Arg	Leu	Leu
				290					295					300
Glu	Tyr	Tyr	Arg	Leu	Tyr	Asn	Thr	Leu	Asp	Asp	Leu	Leu	Leu	Tyr
				305					310					315
Ile	Asn	Ala	Arg	Glu	Leu	Arg	Ile	Thr	Tyr	Gly	Gln	Gly	Ser	Gly
				320					325					330
Thr	Ala	Val	Tyr	Asn	Asn	Asn	Met	Tyr	Val	Asn	Met	Tyr	Asn	Thr
				335					340					345
Gly	Asn	Ile	Ala	Arg	Val	Asn	Leu	Thr	Thr	Asn	Thr	Ile	Ala	Val
				350					355					360
Thr	Gln	Thr	Leu	Pro	Asn	Ala	Ala	Tyr	Asn	Asn	Arg	Phe	Ser	Tyr
				365					370					375
Ala	Asn	Val	Ala	Trp	Gln	Asp	Ile	Asp	Phe	Ala	Val	Asp	Glu	Asn
				380					385					390
Gly	Leu	Trp	Val	Ile	Tyr	Ser	Thr	Glu	Ala	Ser	Thr	Gly	Asn	Met
				395					400					405
Val	Ile	Ser	Lys	Leu	Asn	Asp	Thr	Thr	Leu	Gln	Val	Leu	Asn	Thr
				410					415					420
Trp	Tyr	Thr	Lys	Gln	Tyr	Lys	Pro	Ser	Ala	Ser	Asn	Ala	Phe	Met
				425					430					435
Val	Cys	Gly	Val	Leu	Tyr	Ala	Thr	Arg	Thr	Met	Asn	Thr	Arg	Thr
				440					445					450
Glu	Glu	Ile	Phe	Tyr	Tyr	Tyr	Asp	Thr	Asn	Thr	Gly	Lys	Glu	Gly
				455					460					465
Lys	Leu	Asp	Ile	Val	Met	His	Lys	Met	Gln	Glu	Lys	Val	Gln	Ser
				470					475					480

Ile	Asn	Tyr	Asn	Pro	Phe	Asp	Gln	Lys	Leu	Tyr	Val	Tyr	Asn	Asp
				485					490					495
Gly	Tyr	Leu	Leu	Asn	Tyr	Asp	Leu	Ser	Val	Leu	Gln	Lys	Pro	Gln
				500					505					510

<210> 18  
 <211> 185  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3683905CD1

<400> 18  
 Met Phe Leu Leu Asp Ser Ser Ala Ser Val Ser His Tyr Glu Phe  
 1 5 10 15  
 Ser Arg Val Arg Glu Phe Val Gly Gln Leu Val Ala Pro Leu Pro  
 20 25 30  
 Leu Gly Thr Gly Ala Leu Arg Ala Ser Leu Val His Val Gly Ser  
 35 40 45  
 Arg Pro Tyr Thr Glu Phe Pro Phe Gly Gln His Ser Ser Gly Glu  
 50 55 60  
 Ala Ala Gln Asp Ala Val Arg Ala Ser Ala Gln Arg Met Gly Asp  
 65 70 75  
 Thr His Thr Gly Leu Ala Leu Val Tyr Ala Lys Glu Gln Leu Phe  
 80 85 90  
 Ala Glu Ala Ser Gly Ala Arg Pro Gly Val Pro Lys Val Leu Val  
 95 100 105  
 Trp Val Thr Asp Gly Gly Ser Ser Asp Pro Val Gly Pro Pro Met  
 110 115 120  
 Gln Glu Leu Lys Asp Leu Gly Val Thr Val Phe Ile Val Ser Thr  
 125 130 135  
 Gly Arg Gly Asn Phe Leu Glu Leu Ser Ala Ala Ala Ser Ala Pro  
 140 145 150  
 Ala Glu Lys His Leu His Phe Val Asp Val Asp Asp Leu His Ile  
 155 160 165  
 Ile Val Gln Glu Leu Arg Gly Ser Ile Leu Asp Ala Met Arg Pro  
 170 175 180  
 Gln Ala Tyr Ser Leu  
 185

<210> 19  
 <211> 207  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 4062841CD1

<400> 19  
 Met Ala Ala Leu Val Glu Pro Leu Gly Leu Glu Arg Asp Val Ser  
 1 5 10 15  
 Arg Ala Val Glu Leu Leu Glu Arg Leu Gln Arg Ser Gly Glu Leu  
 20 25 30  
 Pro Pro Gln Lys Leu Gln Ala Leu Gln Arg Val Leu Gln Ser Arg  
 35 40 45  
 Phe Cys Ser Ala Ile Arg Glu Val Tyr Glu Gln Leu Tyr Asp Thr  
 50 55 60  
 Leu Asp Ile Thr Gly Ser Ala Glu Ile Arg Ala His Ala Thr Ala  
 65 70 75  
 Lys Ala Thr Val Ala Ala Phe Thr Ala Ser Glu Gly His Ala His  
 80 85 90  
 Pro Arg Val Val Glu Leu Pro Lys Thr Asp Glu Gly Leu Gly Phe  
 95 100 105

Asn	Ile	Met	Gly	Gly	Lys	Glu	Gln	Asn	Ser	Pro	Ile	Tyr	Ile	Ser	
				110					115					120	
Arg	Val	Ile	Pro	Gly	Gly	Val	Ala	Asp	Arg	His	Gly	Gly	Leu	Lys	
				125					130					135	
Arg	Gly	Asp	Gln	Leu	Leu	Ser	Val	Asn	Gly	Val	Ser	Val	Glu	Gly	
				140					145					150	
Glu	Gln	His	Glu	Lys	Ala	Val	Glu	Leu	Leu	Lys	Ala	Ala	Gln	Gly	
				155					160					165	
Ser	Val	Lys	Leu	Val	Val	Arg	Tyr	Thr	Pro	Arg	Val	Leu	Glu	Glu	
				170					175					180	
Met	Glu	Ala	Arg	Phe	Glu	Lys	Met	Arg	Ser	Ala	Arg	Arg	Arg	Gln	
				185					190					195	
Gln	His	Gln	Ser	Tyr	Ser	Ser	Leu	Glu	Ser	Arg	Gly				
				200					205						

<210> 20  
 <211> 238  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 6394358CD1

<400> 20

Met	Ser	Leu	Asn	Glu	His	Ser	Met	Gln	Ala	Leu	Ser	Trp	Arg	Lys	
1				5					10					15	
Leu	Tyr	Leu	Ser	Arg	Ala	Lys	Leu	Lys	Ala	Ser	Ser	Arg	Thr	Ser	
				20					25					30	
Ala	Leu	Leu	Ser	Gly	Phe	Ala	Met	Val	Ala	Met	Val	Glu	Val	Gln	
				35					40					45	
Leu	Asp	Ala	Asp	His	Asp	Tyr	Pro	Pro	Gly	Leu	Leu	Ile	Ala	Phe	
				50					55					60	
Ser	Ala	Cys	Thr	Thr	Val	Leu	Val	Ala	Val	His	Leu	Phe	Ala	Leu	
				65					70					75	
Met	Ile	Ser	Thr	Cys	Ile	Leu	Pro	Asn	Ile	Glu	Ala	Val	Ser	Asn	
				80					85					90	
Val	His	Asn	Leu	Asn	Ser	Val	Lys	Glu	Ser	Pro	His	Glu	Arg	Met	
				95					100					105	
His	Arg	His	Ile	Glu	Leu	Ala	Trp	Ala	Phe	Ser	Thr	Val	Ile	Gly	
				110					115					120	
Thr	Leu	Leu	Phe	Leu	Ala	Glu	Val	Val	Leu	Leu	Cys	Trp	Val	Lys	
				125					130					135	
Phe	Leu	Pro	Leu	Lys	Lys	Gln	Pro	Gly	Gln	Pro	Arg	Pro	Thr	Ser	
				140					145					150	
Lys	Pro	Pro	Ala	Ser	Gly	Ala	Ala	Ala	Asn	Val	Ser	Thr	Ser	Gly	
				155					160					165	
Ile	Thr	Pro	Gly	Gln	Ala	Ala	Ala	Ile	Ala	Ser	Thr	Thr	Ile	Met	
				170					175					180	
Val	Pro	Phe	Gly	Leu	Ile	Phe	Ile	Val	Phe	Ala	Val	His	Phe	Tyr	
				185					190					195	
Arg	Ser	Leu	Val	Ser	His	Lys	Thr	Asp	Arg	Gln	Phe	Gln	Glu	Leu	
				200					205					210	
Asn	Glu	Leu	Ala	Glu	Phe	Ala	Arg	Leu	Gln	Asp	Gln	Leu	Asp	His	
				215					220					225	
Arg	Gly	Asp	His	Pro	Leu	Thr	Pro	Gly	Ser	His	Tyr	Ala			
				230					235						

<210> 21  
 <211> 3298  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2847752CD1

&lt;400&gt; 21

```

Met Ala Arg Arg Pro Pro Trp Arg Gly Leu Gly Gly Arg Ser Thr
1      5      10      15
Pro Ile Leu Leu Leu Leu Leu Leu Ser Leu Phe Pro Leu Ser Gln
20     25     30
Glu Glu Leu Gly Gly Gly Gly His Gln Gly Trp Asp Pro Gly Leu
35     40     45
Ala Ala Thr Thr Gly Pro Arg Ala His Ile Gly Gly Gly Ala Leu
50     55     60
Ala Leu Cys Pro Glu Ser Ser Gly Val Arg Glu Asp Gly Gly Pro
65     70     75
Gly Leu Gly Val Arg Glu Pro Ile Phe Val Gly Leu Arg Gly Arg
80     85     90
Arg Gln Ser Ala Arg Asn Ser Arg Gly Pro Pro Glu Gln Pro Asn
95     100    105
Glu Glu Leu Gly Ile Glu His Gly Val Gln Pro Leu Gly Ser Arg
110    115    120
Glu Arg Glu Thr Gly Gln Gly Pro Gly Ser Val Leu Tyr Trp Arg
125    130    135
Pro Glu Val Ser Ser Cys Gly Arg Thr Gly Pro Leu Gln Arg Gly
140    145    150
Ser Leu Ser Pro Gly Ala Leu Ser Ser Gly Val Pro Gly Ser Gly
155    160    165
Asn Ser Ser Pro Leu Pro Ser Asp Phe Leu Ile Arg His His Gly
170    175    180
Pro Lys Pro Val Ser Ser Gln Arg Asn Ala Gly Thr Gly Ser Arg
185    190    195
Lys Arg Val Gly Thr Ala Arg Cys Cys Gly Glu Leu Trp Ala Thr
200    205    210
Gly Ser Lys Gly Gln Gly Glu Arg Ala Thr Thr Ser Gly Ala Glu
215    220    225
Arg Thr Ala Pro Arg Arg Asn Cys Leu Pro Gly Ala Ser Gly Ser
230    235    240
Gly Pro Glu Leu Asp Ser Ala Pro Arg Thr Ala Arg Thr Ala Pro
245    250    255
Ala Ser Gly Ser Ala Pro Arg Glu Ser Arg Thr Ala Pro Glu Pro
260    265    270
Ala Pro Lys Arg Met Arg Ser Arg Gly Leu Phe Arg Cys Arg Phe
275    280    285
Leu Pro Gln Arg Pro Gly Pro Arg Pro Pro Gly Leu Pro Ala Arg
290    295    300
Pro Glu Ala Arg Lys Val Thr Ser Ala Asn Arg Ala Arg Phe Arg
305    310    315
Arg Ala Ala Asn Arg His Pro Gln Phe Pro Gln Tyr Asn Tyr Gln
320    325    330
Thr Leu Val Pro Glu Asn Glu Ala Ala Gly Thr Ala Val Leu Arg
335    340    345
Val Val Ala Gln Asp Pro Asp Ala Gly Glu Ala Gly Arg Leu Val
350    355    360
Tyr Ser Leu Ala Ala Leu Met Asn Ser Arg Ser Leu Glu Leu Phe
365    370    375
Ser Ile Asp Pro Gln Ser Gly Leu Ile Arg Thr Ala Ala Ala Leu
380    385    390
Asp Arg Glu Ser Met Glu Arg His Tyr Leu Arg Val Thr Ala Gln
395    400    405
Asp His Gly Ser Pro Arg Leu Ser Ala Thr Thr Met Val Ala Val
410    415    420
Thr Val Ala Asp Arg Asn Asp His Ser Pro Val Phe Glu Gln Ala
425    430    435
Gln Tyr Arg Glu Thr Leu Arg Glu Asn Val Glu Glu Gly Tyr Pro
440    445    450
Ile Leu Gln Leu Arg Ala Thr Asp Gly Asp Ala Pro Pro Asn Ala
455    460    465
Asn Leu Arg Tyr Arg Phe Val Gly Pro Pro Ala Ala Arg Ala Ala
470    475    480
Ala Ala Ala Ala Phe Glu Ile Asp Pro Arg Ser Gly Leu Ile Ser
485    490    495

```

Thr	Ser	Gly	Arg	Val	Asp	Arg	Glu	His	Met	Glu	Ser	Tyr	Glu	Leu
				500					505					510
Val	Val	Glu	Ala	Ser	Asp	Gln	Gly	Gln	Glu	Pro	Gly	Pro	Arg	Ser
				515					520					525
Ala	Thr	Val	Arg	Val	His	Ile	Thr	Val	Leu	Asp	Glu	Asn	Asp	Asn
				530					535					540
Ala	Pro	Gln	Phe	Ser	Glu	Lys	Arg	Tyr	Val	Ala	Gln	Val	Arg	Glu
				545					550					555
Asp	Val	Arg	Pro	His	Thr	Val	Val	Leu	Arg	Val	Thr	Ala	Thr	Asp
				560					565					570
Arg	Asp	Lys	Asp	Ala	Asn	Gly	Leu	Val	His	Tyr	Asn	Ile	Ile	Ser
				575					580					585
Gly	Asn	Ser	Arg	Gly	His	Phe	Ala	Ile	Asp	Ser	Leu	Thr	Gly	Glu
				590					595					600
Ile	Gln	Val	Val	Ala	Pro	Leu	Asp	Phe	Glu	Ala	Glu	Arg	Glu	Tyr
				605					610					615
Ala	Leu	Arg	Ile	Arg	Ala	Gln	Asp	Ala	Gly	Arg	Pro	Pro	Leu	Ser
				620					625					630
Asn	Asn	Thr	Gly	Leu	Ala	Ser	Ile	Gln	Val	Val	Asp	Ile	Asn	Asp
				635					640					645
His	Ile	Pro	Ile	Phe	Val	Ser	Thr	Pro	Phe	Gln	Val	Ser	Val	Leu
				650					655					660
Glu	Asn	Ala	Pro	Leu	Gly	His	Ser	Val	Ile	His	Ile	Gln	Ala	Val
				665					670					675
Asp	Ala	Asp	His	Gly	Glu	Asn	Ala	Arg	Leu	Glu	Tyr	Ser	Leu	Thr
				680					685					690
Gly	Val	Ala	Pro	Asp	Thr	Pro	Phe	Val	Ile	Asn	Ser	Ala	Thr	Gly
				695					700					705
Trp	Val	Ser	Val	Ser	Gly	Pro	Leu	Asp	Arg	Glu	Ser	Val	Glu	His
				710					715					720
Tyr	Phe	Phe	Gly	Val	Glu	Ala	Arg	Asp	His	Gly	Ser	Pro	Pro	Leu
				725					730					735
Ser	Ala	Ser	Ala	Ser	Val	Thr	Val	Thr	Val	Leu	Asp	Val	Asn	Asp
				740					745					750
Asn	Arg	Pro	Glu	Phe	Thr	Met	Lys	Glu	Tyr	His	Leu	Arg	Leu	Asn
				755					760					765
Glu	Asp	Ala	Ala	Val	Gly	Thr	Ser	Val	Val	Ser	Val	Thr	Ala	Val
				770					775					780
Asp	Arg	Asp	Ala	Asn	Ser	Ala	Ile	Ser	Tyr	Gln	Ile	Thr	Gly	Gly
				785					790					795
Asn	Thr	Arg	Asn	Arg	Phe	Ala	Ile	Ser	Thr	Gln	Gly	Gly	Val	Gly
				800					805					810
Leu	Val	Thr	Leu	Ala	Leu	Pro	Leu	Asp	Tyr	Lys	Gln	Glu	Arg	Tyr
				815					820					825
Phe	Lys	Leu	Val	Leu	Thr	Ala	Ser	Asp	Arg	Ala	Leu	His	Asp	His
				830					835					840
Cys	Tyr	Val	His	Ile	Asn	Ile	Thr	Asp	Ala	Asn	Thr	His	Arg	Pro
				845					850					855
Val	Phe	Gln	Ser	Ala	His	Tyr	Ser	Val	Ser	Val	Asn	Glu	Asp	Arg
				860					865					870
Pro	Met	Gly	Ser	Thr	Ile	Val	Val	Ile	Ser	Ala	Ser	Asp	Asp	Asp
				875					880					885
Val	Gly	Glu	Asn	Ala	Arg	Ile	Thr	Tyr	Leu	Leu	Glu	Asp	Asn	Leu
				890					895					900
Pro	Gln	Phe	Arg	Ile	Asp	Ala	Asp	Ser	Gly	Ala	Ile	Thr	Leu	Gln
				905					910					915
Ala	Pro	Leu	Asp	Tyr	Glu	Asp	Gln	Val	Thr	Tyr	Thr	Leu	Ala	Ile
				920					925					930
Thr	Ala	Arg	Asp	Asn	Gly	Ile	Pro	Gln	Lys	Ala	Asp	Thr	Thr	Tyr
				935					940					945
Val	Glu	Val	Met	Val	Asn	Asp	Val	Asn	Asp	Asn	Ala	Pro	Gln	Phe
				950					955					960
Val	Ala	Ser	His	Tyr	Thr	Gly	Leu	Val	Ser	Glu	Asp	Ala	Pro	Pro
				965					970					975
Phe	Thr	Ser	Val	Leu	Gln	Ile	Ser	Ala	Thr	Asp	Arg	Asp	Ala	His
				980					985					990
Ala	Asn	Gly	Arg	Val	Gln	Tyr	Thr	Phe	Gln	Asn	Gly	Glu	Asp	Gly

995	1000	1005
Asp Gly Asp Phe Thr Ile Glu Pro Thr Ser Gly Ile Val Arg Thr		
1010	1015	1020
Val Arg Arg Leu Asp Arg Glu Ala Val Ser Val Tyr Glu Leu Thr		
1025	1030	1035
Ala Tyr Ala Val Asp Arg Gly Val Pro Pro Leu Arg Thr Pro Val		
1040	1045	1050
Ser Ile Gln Val Met Val Gln Asp Val Asn Asp Asn Ala Pro Val		
1055	1060	1065
Phe Pro Ala Glu Glu Phe Glu Val Arg Val Lys Glu Asn Ser Ile		
1070	1075	1080
Val Gly Ser Val Val Ala Gln Ile Thr Ala Val Asp Pro Asp Glu		
1085	1090	1095
Gly Pro Asn Ala His Ile Met Tyr Gln Ile Val Glu Gly Asn Ile		
1100	1105	1110
Pro Glu Leu Phe Gln Met Asp Ile Phe Ser Gly Glu Leu Thr Ala		
1115	1120	1125
Leu Ile Asp Leu Asp Tyr Glu Ala Arg Gln Glu Tyr Val Ile Val		
1130	1135	1140
Val Gln Ala Thr Ser Ala Pro Leu Val Ser Arg Ala Thr Val His		
1145	1150	1155
Val Arg Leu Val Asp Gln Asn Asp Asn Ser Pro Val Leu Asn Asn		
1160	1165	1170
Phe Gln Ile Leu Phe Asn Asn Tyr Val Ser Asn Arg Ser Asp Thr		
1175	1180	1185
Phe Pro Ser Gly Ile Ile Gly Arg Ile Pro Ala Tyr Asp Pro Asp		
1190	1195	1200
Val Ser Asp His Leu Phe Tyr Ser Phe Glu Arg Gly Asn Glu Leu		
1205	1210	1215
Gln Leu Leu Val Val Asn Gln Thr Ser Gly Glu Leu Arg Leu Ser		
1220	1225	1230
Arg Lys Leu Asp Asn Asn Arg Pro Leu Val Ala Ser Met Leu Val		
1235	1240	1245
Thr Val Thr Asp Gly Leu His Ser Val Thr Ala Gln Cys Val Leu		
1250	1255	1260
Arg Val Val Ile Ile Thr Glu Glu Leu Leu Ala Asn Ser Leu Thr		
1265	1270	1275
Val Arg Leu Glu Asn Met Trp Gln Glu Arg Phe Leu Ser Pro Leu		
1280	1285	1290
Leu Gly Arg Phe Leu Glu Gly Val Ala Ala Val Leu Ala Thr Pro		
1295	1300	1305
Ala Glu Asp Val Phe Ile Phe Asn Ile Gln Asn Asp Thr Asp Val		
1310	1315	1320
Gly Gly Thr Val Leu Asn Val Ser Phe Ser Ala Leu Ala Pro Arg		
1325	1330	1335
Gly Ala Gly Ala Gly Ala Ala Gly Pro Trp Phe Ser Ser Glu Glu		
1340	1345	1350
Leu Gln Glu Gln Leu Tyr Val Arg Arg Ala Ala Leu Ala Ala Arg		
1355	1360	1365
Ser Leu Leu Asp Val Leu Pro Phe Asp Asp Asn Val Cys Leu Arg		
1370	1375	1380
Glu Pro Cys Glu Asn Tyr Met Lys Cys Val Ser Val Leu Arg Phe		
1385	1390	1395
Asp Ser Ser Ala Pro Phe Leu Ala Ser Ala Ser Thr Leu Phe Arg		
1400	1405	1410
Pro Ile Gln Pro Ile Ala Gly Leu Arg Cys Arg Cys Pro Pro Gly		
1415	1420	1425
Phe Thr Gly Asp Phe Cys Glu Thr Glu Leu Asp Leu Cys Tyr Ser		
1430	1435	1440
Asn Pro Cys Arg Asn Gly Gly Ala Cys Ala Arg Arg Glu Gly Gly		
1445	1450	1455
Tyr Thr Cys Val Cys Arg Pro Arg Phe Thr Gly Glu Asp Cys Glu		
1460	1465	1470
Leu Asp Thr Glu Ala Gly Arg Cys Val Pro Gly Val Cys Arg Asn		
1475	1480	1485
Gly Gly Thr Cys Thr Asp Ala Pro Asn Gly Gly Phe Arg Cys Gln		
1490	1495	1500

Cys	Pro	Ala	Gly	Gly	Ala	Phe	Glu	Gly	Pro	Arg	Cys	Glu	Val	Ala	1505	1510	1515
Ala	Arg	Ser	Phe	Pro	Pro	Ser	Ser	Phe	Val	Met	Phe	Arg	Gly	Leu	1520	1525	1530
Arg	Gln	Arg	Phe	His	Leu	Thr	Leu	Ser	Leu	Ser	Phe	Ala	Thr	Val	1535	1540	1545
Gln	Gln	Ser	Gly	Leu	Leu	Phe	Tyr	Asn	Gly	Arg	Leu	Asn	Glu	Lys	1550	1555	1560
His	Asp	Phe	Leu	Ala	Leu	Glu	Leu	Val	Ala	Gly	Gln	Val	Arg	Leu	1565	1570	1575
Thr	Tyr	Ser	Thr	Gly	Glu	Ser	Asn	Thr	Val	Val	Ser	Pro	Thr	Val	1580	1585	1590
Pro	Gly	Gly	Leu	Ser	Asp	Gly	Gln	Trp	His	Thr	Val	His	Leu	Arg	1595	1600	1605
Tyr	Tyr	Asn	Lys	Pro	Arg	Thr	Asp	Ala	Leu	Gly	Gly	Ala	Gln	Gly	1610	1615	1620
Pro	Ser	Lys	Asp	Lys	Val	Ala	Val	Leu	Ser	Val	Asp	Asp	Cys	Asp	1625	1630	1635
Val	Ala	Val	Ala	Leu	Gln	Phe	Gly	Ala	Glu	Ile	Gly	Asn	Tyr	Ser	1640	1645	1650
Cys	Ala	Ala	Ala	Gly	Val	Gln	Thr	Ser	Ser	Lys	Lys	Ser	Leu	Asp	1655	1660	1665
Leu	Thr	Gly	Pro	Leu	Leu	Leu	Gly	Gly	Val	Pro	Asn	Leu	Pro	Glu	1670	1675	1680
Asn	Phe	Pro	Val	Ser	His	Lys	Asp	Phe	Ile	Gly	Cys	Met	Arg	Asp	1685	1690	1695
Leu	His	Ile	Asp	Gly	Arg	Arg	Val	Asp	Met	Ala	Ala	Phe	Val	Ala	1700	1705	1710
Asn	Asn	Gly	Thr	Met	Ala	Gly	Cys	Gln	Ala	Lys	Leu	His	Phe	Cys	1715	1720	1725
Asp	Ser	Gly	Pro	Cys	Lys	Asn	Ser	Gly	Phe	Cys	Ser	Glu	Arg	Trp	1730	1735	1740
Gly	Ser	Phe	Ser	Cys	Asp	Cys	Pro	Val	Gly	Phe	Gly	Gly	Lys	Asp	1745	1750	1755
Cys	Gln	Leu	Thr	Met	Ala	His	Pro	His	His	Phe	Arg	Gly	Asn	Gly	1760	1765	1770
Thr	Leu	Ser	Trp	Asn	Phe	Gly	Ser	Asp	Met	Ala	Val	Ser	Val	Pro	1775	1780	1785
Trp	Tyr	Leu	Gly	Leu	Ala	Phe	Arg	Thr	Arg	Ala	Thr	Gln	Gly	Val	1790	1795	1800
Leu	Met	Gln	Val	Gln	Ala	Gly	Pro	His	Ser	Thr	Leu	Leu	Cys	Gln	1805	1810	1815
Leu	Asp	Arg	Gly	Leu	Leu	Ser	Val	Thr	Val	Thr	Arg	Gly	Ser	Gly	1820	1825	1830
Arg	Ala	Ser	His	Leu	Leu	Leu	Asp	Gln	Val	Thr	Val	Ser	Asp	Gly	1835	1840	1845
Arg	Trp	His	Asp	Leu	Arg	Leu	Glu	Leu	Gln	Glu	Glu	Pro	Gly	Gly	1850	1855	1860
Arg	Arg	Gly	His	His	Val	Leu	Met	Val	Ser	Leu	Asp	Phe	Ser	Leu	1865	1870	1875
Phe	Gln	Asp	Thr	Met	Ala	Val	Gly	Ser	Glu	Leu	Gln	Gly	Leu	Lys	1880	1885	1890
Val	Lys	Gln	Leu	His	Val	Gly	Gly	Leu	Pro	Pro	Gly	Ser	Ala	Glu	1895	1900	1905
Glu	Ala	Pro	Gln	Gly	Leu	Val	Gly	Cys	Ile	Gln	Pro	Pro	Ser	Glu	1910	1915	1920
Cys	Gly	Pro	Gly	Cys	Val	Val	Thr	Asn	Ala	Cys	Ala	Ser	Gly	Pro	1925	1930	1935
Cys	Pro	Pro	His	Ala	Asp	Cys	Arg	Asp	Leu	Trp	Gln	Thr	Phe	Ser	1940	1945	1950
Cys	Thr	Cys	Gln	Pro	Gly	Tyr	Tyr	Gly	Pro	Gly	Cys	Val	Asp	Ala	1955	1960	1965
Cys	Leu	Leu	Asn	Pro	Cys	Gln	Asn	Gln	Gly	Ser	Cys	Arg	His	Leu	1970	1975	1980
Pro	Gly	Ala	Pro	His	Gly	Tyr	Thr	Cys	Asp	Cys	Val	Gly	Gly	Tyr	1985	1990	1995
Phe	Gly	His	His	Cys	Glu	His	Arg	Met	Asp	Gln	Gln	Cys	Pro	Arg			



2000	2005	2010
Gly Trp Trp Gly Ser	Pro Thr Cys Gly Pro	Cys Asn Cys Asp Val
2015	2020	2025
His Lys Gly Phe Asp	Pro Asn Cys Asn Lys	Thr Asn Gly Gln Cys
2030	2035	2040
His Cys Lys Glu Phe	His Tyr Arg Pro Arg	Gly Ser Asp Ser Cys
2045	2050	2055
Leu Pro Cys Asp Cys	Tyr Pro Val Gly Ser	Thr Ser Arg Ser Cys
2060	2065	2070
Ala Pro His Ser Gly	Gln Cys Pro Cys Arg	Pro Gly Ala Leu Gly
2075	2080	2085
Arg Gln Cys Asn Ser	Cys Asp Ser Pro Phe	Ala Glu Val Thr Ala
2090	2095	2100
Ser Gly Cys Arg Val	Leu Tyr Asp Ala Cys	Pro Lys Ser Leu Arg
2105	2110	2115
Ser Gly Val Trp Trp	Pro Gln Thr Lys Phe	Gly Val Leu Ala Thr
2120	2125	2130
Val Pro Cys Pro Arg	Gly Ala Leu Gly Leu	Arg Gly Ala Gly Ala
2135	2140	2145
Ala Val Arg Leu Cys	Asp Glu Ala Gln Gly	Trp Leu Glu Pro Asp
2150	2155	2160
Leu Phe Asn Cys Thr	Ser Pro Ala Phe Arg	Glu Leu Ser Leu Leu
2165	2170	2175
Leu Asp Gly Leu Glu	Leu Asn Lys Thr Ala	Leu Asp Thr Met Glu
2180	2185	2190
Ala Lys Lys Leu Ala	Gln Arg Leu Arg Glu	Val Thr Gly His Thr
2195	2200	2205
Asp His Tyr Phe Ser	Gln Asp Val Arg Val	Thr Ala Arg Leu Leu
2210	2215	2220
Ala His Leu Leu Ala	Phe Glu Ser His Gln	Gln Gly Phe Gly Leu
2225	2230	2235
Thr Ala Thr Gln Asp	Ala His Phe Asn Glu	Asn Leu Leu Trp Ala
2240	2245	2250
Gly Ser Ala Leu Leu	Ala Pro Glu Thr Gly	Asp Leu Trp Ala Ala
2255	2260	2265
Leu Gly Gln Arg Ala	Pro Gly Gly Ser Pro	Gly Ser Ala Gly Leu
2270	2275	2280
Val Arg His Leu Glu	Glu Tyr Ala Ala Thr	Leu Ala Arg Asn Met
2285	2290	2295
Glu Leu Thr Tyr Leu	Asn Pro Met Gly Leu	Val Thr Pro Asn Ile
2300	2305	2310
Met Leu Ser Ile Asp	Arg Met Glu His Pro	Ser Ser Pro Arg Gly
2315	2320	2325
Ala Arg Arg Tyr Pro	Arg Tyr His Ser Asn	Leu Phe Arg Gly Gln
2330	2335	2340
Asp Ala Trp Asp Pro	His Thr His Val Leu	Leu Pro Ser Gln Ser
2345	2350	2355
Pro Arg Pro Ser Pro	Ser Glu Val Leu Pro	Thr Ser Ser Ser Ile
2360	2365	2370
Glu Asn Ser Thr Thr	Ser Ser Val Val Pro	Pro Pro Ala Pro Pro
2375	2380	2385
Glu Pro Glu Pro Gly	Ile Ser Ile Ile Ile	Leu Leu Val Tyr Arg
2390	2395	2400
Thr Leu Gly Gly Leu	Leu Pro Ala Gln Phe	Gln Ala Glu Arg Arg
2405	2410	2415
Gly Ala Arg Leu Pro	Gln Asn Pro Val Met	Asn Ser Pro Val Val
2420	2425	2430
Ser Val Ala Val Phe	His Gly Arg Asn Phe	Leu Arg Gly Ile Leu
2435	2440	2445
Glu Ser Pro Ile Ser	Leu Glu Phe Arg Leu	Leu Gln Thr Ala Asn
2450	2455	2460
Arg Ser Lys Ala Ile	Cys Val Gln Trp Asp	Pro Pro Gly Leu Ala
2465	2470	2475
Glu Gln His Gly Val	Trp Thr Ala Arg Asp	Cys Glu Leu Val His
2480	2485	2490
Arg Asn Gly Ser His	Ala Arg Cys Arg Cys	Ser Arg Thr Gly Thr
2495	2500	2505

Phe Gly Val Leu Met	Asp Ala Ser Pro Arg	Glu Arg Leu Glu Gly	2510	2515	2520
Asp Leu Glu Leu Leu	Ala Val Phe Thr His	Val Val Val Ala Val	2525	2530	2535
Ser Val Ala Ala Leu	Val Leu Thr Ala Ala	Ile Leu Leu Ser Leu	2540	2545	2550
Arg Ser Leu Lys Ser	Asn Val Arg Gly Ile	His Ala Asn Val Ala	2555	2560	2565
Ala Ala Leu Gly Val	Ala Glu Leu Leu Phe	Leu Leu Gly Ile His	2570	2575	2580
Arg Thr His Asn Gln	Leu Val Cys Thr Ala	Val Ala Ile Leu Leu	2585	2590	2595
His Tyr Phe Phe Leu	Ser Thr Phe Ala Trp	Leu Phe Val Gln Gly	2600	2605	2610
Leu His Leu Tyr Arg	Met Gln Val Glu Pro	Arg Asn Val Asp Arg	2615	2620	2625
Gly Ala Met Arg Phe	Tyr His Ala Leu Gly	Trp Gly Val Pro Ala	2630	2635	2640
Val Leu Leu Gly Leu	Ala Val Gly Leu Asp	Pro Glu Gly Tyr Gly	2645	2650	2655
Asn Pro Asp Phe Cys	Trp Ile Ser Val His	Glu Pro Leu Ile Trp	2660	2665	2670
Ser Phe Ala Gly Pro	Val Val Leu Val Ile	Val Met Asn Gly Thr	2675	2680	2685
Met Phe Leu Leu Ala	Ala Arg Thr Ser Cys	Ser Thr Gly Gln Arg	2690	2695	2700
Glu Ala Lys Lys Thr	Ser Ala Leu Thr Leu	Arg Ser Ser Phe Leu	2705	2710	2715
Leu Leu Leu Leu Val	Ser Ala Ser Trp Leu	Phe Gly Leu Leu Ala	2720	2725	2730
Val Asn His Ser Ile	Leu Ala Phe His Tyr	Leu His Ala Gly Leu	2735	2740	2745
Cys Gly Leu Gln Gly	Leu Ala Val Leu Leu	Leu Phe Cys Val Leu	2750	2755	2760
Asn Ala Asp Ala Arg	Ala Ala Trp Met Pro	Ala Cys Leu Gly Arg	2765	2770	2775
Lys Ala Ala Pro Glu	Glu Ala Arg Pro Ala	Pro Gly Leu Gly Pro	2780	2785	2790
Gly Ala Tyr Asn Asn	Thr Ala Leu Phe Glu	Glu Ser Gly Leu Ile	2795	2800	2805
Arg Ile Thr Leu Gly	Ala Ser Thr Val Ser	Ser Val Ser Ser Ala	2810	2815	2820
Arg Ser Gly Arg Thr	Gln Asp Gln Asp Ser	Gln Arg Gly Arg Ser	2825	2830	2835
Tyr Leu Arg Asp Asp	Val Leu Val Arg His	Gly Ser Ala Ala Asp	2840	2845	2850
His Thr Asp His Ser	Leu Gln Ala His Ala	Gly Pro Thr Asp Leu	2855	2860	2865
Asp Val Ala Met Phe	His Arg Asp Ala Gly	Ala Asp Ser Asp Ser	2870	2875	2880
Asp Ser Asp Leu Ser	Leu Glu Glu Glu Arg	Ser Leu Ser Ile Pro	2885	2890	2895
Ser Ser Glu Ser Glu	Asp Asn Gly Arg Thr	Arg Gly Arg Val Gln	2900	2905	2910
Arg Pro Leu Cys Arg	Ala Ala Gln Ser Glu	Arg Leu Leu Thr His	2915	2920	2925
Pro Lys Asp Val Asp	Gly Asn Asp Leu Leu	Ser Tyr Trp Pro Ala	2930	2935	2940
Leu Gly Glu Cys Glu	Ala Ala Pro Cys Ala	Leu Gln Thr Trp Gly	2945	2950	2955
Ser Glu Arg Arg Leu	Gly Leu Asp Thr Ser	Lys Asp Ala Ala Asn	2960	2965	2970
Asn Asn Gln Pro Asp	Pro Ala Leu Thr Ser	Gly Asp Glu Thr Ser	2975	2980	2985
Leu Gly Arg Ala Gln	Arg Gln Arg Lys Gly	Ile Leu Lys Asn Arg	2990	2995	3000
Leu Gln Tyr Pro Leu	Val Pro Gln Thr Arg	Gly Ala Pro Glu Leu			

3005	3010	3015
Ser Trp Cys Arg Ala	Ala Thr Leu Gly His Arg Ala Val Pro Ala	
3020	3025	3030
Ala Ser Tyr Gly Arg	Ile Tyr Ala Gly Gly Gly Thr Gly Ser Leu	
3035	3040	3045
Ser Gln Pro Ala Ser	Arg Tyr Ser Ser Arg Glu Gln Leu Asp Leu	
3050	3055	3060
Leu Leu Arg Arg Gln	Leu Ser Arg Glu Arg Leu Glu Glu Ala Pro	
3065	3070	3075
Ala Pro Val Leu Arg	Pro Leu Ser Arg Pro Gly Ser Gln Glu Cys	
3080	3085	3090
Met Asp Ala Ala Pro	Gly Arg Leu Glu Pro Lys Asp Arg Gly Ser	
3095	3100	3105
Thr Leu Pro Arg Arg	Gln Pro Pro Arg Asp Tyr Pro Gly Ala Met	
3110	3115	3120
Ala Gly Arg Phe Gly	Ser Arg Asp Ala Leu Asp Leu Gly Ala Pro	
3125	3130	3135
Arg Glu Trp Leu Ser	Thr Leu Pro Pro Pro Arg Arg Thr Arg Asp	
3140	3145	3150
Leu Asp Pro Gln Pro	Pro Pro Leu Pro Leu Ser Pro Gln Arg Gln	
3155	3160	3165
Leu Ser Arg Asp Pro	Leu Leu Pro Ser Arg Pro Leu Asp Ser Leu	
3170	3175	3180
Ser Arg Ser Ser Asn	Ser Arg Glu Gln Leu Asp Gln Val Pro Ser	
3185	3190	3195
Arg His Pro Ser Arg	Glu Ala Leu Gly Pro Leu Pro Gln Leu Leu	
3200	3205	3210
Arg Ala Arg Glu Asp	Ser Val Ser Gly Pro Ser His Gly Pro Ser	
3215	3220	3225
Thr Glu Gln Leu Asp	Ile Leu Ser Ser Ile Leu Ala Ser Phe Asn	
3230	3235	3240
Ser Ser Ala Leu Ser	Ser Val Gln Ser Ser Ser Thr Pro Leu Gly	
3245	3250	3255
Pro His Thr Thr Ala	Thr Pro Ser Ala Thr Ala Ser Val Leu Gly	
3260	3265	3270
Pro Ser Thr Pro Arg	Ser Ala Thr Ser His Ser Ile Ser Glu Leu	
3275	3280	3285
Ser Pro Asp Ser Glu	Val Pro Arg Ser Glu Gly His Ser	
3290	3295	

<210> 22  
 <211> 905  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 1424691CB1

<400> 22  
 cccacgcgtc cgtggttagaa ctccaaaaag cactaaacat gtccattcga ttcagtggag 60  
 tagaacaaaa cctcaggatg aagtgaagc agtccaactt gccattcaga cattattcac 120  
 caattcagat ggcaaccctg gaagcaggct cgactcaagt gctgattgcc agtgggttaga 180  
 tactctgagg atgcggcaga ttgcatccaa cacttcttta cagcgttccc agagcaatcc 240  
 tattctgggg tcaccgttct tctcacactt tgatggccag gattcctacg ctgctgctgt 300  
 gagacggccc caggtgcccc ttaagtatca acagattaca cctgtgaacc agtcacagaag 360  
 ctgctctcct actcagtatg gactgaccaa aaacttctct tccctacatc tcaactctag 420  
 ggacagtggc ttttccagtg gcaatactga cacctcttca gagaggggtc gatactcaga 480  
 cagaagcagg aacaaatatg gacgtggtag tatatcactc aattcttctc ctgaggaag 540  
 atacagtgga aagagtcagc attccactcc ttcaagagga agataccctg gaaagtctta 600  
 cagggtttct cagtcagcac tcaatcctca ccagtcgcct gacttcaaga gaagccccag 660  
 ggacctccac caacccaaca ccataaccagg gatgcctttg caccctgaga ctgactcaag 720  
 agccagtgaag gaggcagca aagtcagcga agggggctgg acaaaagtgg aataccggaa 780  
 aaagccccac aggccatctc ccgccaacac caataaagag agagccagag gggaccaccg 840  
 tggatggaga aacttttgat gaattgaact acatagcttt tctaagcagg ttaaaaaaaa 900  
 aaaaaa 905

<210> 23  
<211> 2390  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 1450801CB1

<400> 23  
tgaccttggc accccgcttc ctgccaaacg ccctgatcag tgtggacttt gctgccaact 60  
atctcaccaa gatctatggg ctcacctttg gccagaagcc aaacttgagg tctgtgtacc 120  
tgcacaacaa caagctggca gacgcccggc tgccggacaa catgttcaac ggctccagca 180  
acgtcgagggt cctcatcctg tccagcaact tcttgcccca cgtgcccagg cacctgccgc 240  
ctgccctgta caagctgcac ctcaagaaca acaagctgga gaagatcccc ccggggggcct 300  
tgcagcgagc tggagcagcc tgcgcgagct atacctgcag aacaactacc tgactgacga 360  
gggcctggac aacgagacct tctggaagct ctccagcctg gagtacctgg atctgtccag 420  
caacaacctg tctcgggtcc cagctgggct gccgcgcagc ctgggtgctgc tgcacttgga 480  
gaagaacgcc atccggagcg tggacgcgaa tgtgctgacc cccatccgca gcctggagta 540  
cctgctgctg cacagcaacc agctgcggga gcagggcac caccactgg ccttccaggg 600  
cctcaaggcg ttgcacacgg tgcacctgta caacaacgcy ctggagcgcg tgcccagtgg 660  
cctgcctcgc cgcgtgcgca cctcatgat cctgcacaac cagatcacag gcattggccg 720  
cgaagacttt gccaccacct acttctctgga ggagctcaac ctgagctaca accgcatcac 780  
cagcccacag gtgcaccgcy acgcttcccg caagctgcgc ctgctgcgct cgctggacct 840  
gtcgggcaac cggctgcaca cgctgccacc tgggctgcct cgaaatgtcc atgtgctgaa 900  
ggtcgaagcg aatgagctgc ctgccttggc acgaggggcy ctggcgggca tggtcagct 960  
gcgtgagctg tacctacca gcaaccgact gcgcagccga gccctgggcy cccgtgcctg 1020  
gggtggacctc gcccatctgc agctgctgga catcgccggg aatcagctca cagagatccc 1080  
cgaggggctc cccgagtcac ttgagtacct gtacctgcag aacaacaaga ttagtgcggt 1140  
gcccgccaat gccttcgact ccacgcccaa cctcaagggg atctttctca ggtttaacaa 1200  
gctggctgtg ggctccgtgg tggacagtgc ctcccgagg ctgaagcacc tgcaggctct 1260  
ggacattgaa ggcaacttag agtttgggtga catttccaag gaccgtggcc gcttggggaa 1320  
ggaaaaggag gaggaggaag aggaggagga ggaggaagag gaaacaagat agtgacaagg 1380  
tgatgcagat gtgacctagg atgatggacc gccggactct tttctgcagc acacgcctgt 1440  
gtgctgtgag ccccccactc tgccgtgctc acacagacac acccagctgc acacatgagg 1500  
catcccatcat gacacgggct gacacagtct catatcccca ccccttccca cggcgtgtcc 1560  
cacggccaga cacatgcaca cacatcacac cctcaaacac ccagctcagc cacacacaac 1620  
tacctccaa accaccacag tctctgtcac acccccacta ccgctgccac gccctctgaa 1680  
tcatgcaggg aagggctgc ccctgccctg gcacacacag gcacccattc cctccccctg 1740  
ctgacatgtg tatgcgtatg catacacacc acacacacac acatgcacaa gtcattgtcg 1800  
aacagccctc caaagcctat gccacagaca gctcttgccc cagccagaat cagccatagc 1860  
agctcgccgt ctgccctgtc catctgtccg tccgttccct ggagaagaca caagggatc 1920  
catgctctgt ggccagggtc ctgccacctc ctggaactca caaaagctgg cttttattcc 1980  
tttcccatcc tatggggaca ggagccttca ggactgctgg cctggcctgg cccaccctgc 2040  
tctccagggt gctgggcagt cactctgcta agagtccctc cctggccagc cctggcagga 2100  
cacaggcaact tttccaatgg gcaagcccag tggaggcagg atgggagagc cccctgggtg 2160  
ctgctggggc cttggggcag gagtgaagca gaggtgatgg ggctgggctg agccagggag 2220  
gaaggaccca gctgcaccta ggagacacct ttgttcttca ggctgtggg ggaagtccg 2280  
ggtgccttta tttttattc ttttctaagg aaaaaaatga taaaaatctc aaagctgatt 2340  
tttcttgtaa tagaaaaact aatataaaag cattatccct aaaaaaaaaa 2390

<210> 24  
<211> 1471  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 1597872CB1

<400> 24  
cttggctagg agcccgtgc gggccaggcg cgcggtgcgc aggcataccc gggctcgccgg 60  
tgaacaccgc ggagccccgg gacttgccgt tcgttgctgg tgccggcatg gtcttctgtc 120  
tgtcgagcga ggagccgcgc cgcccgtgc gaacgacatg gtccacttcc aggcctcgga 180  
agtccagcag ctgctacaca acaagtctgt ggtcatcttg ggggactcca ttcagagggc 240  
tgtgtacaag gacctggtgc tcttgetcca gaaagactca ctgctcacag ctgccagct 300  
gaaagccaag taccttgagg atgttctgga agagctgaca tatggacctg ccccgacct 360

```

ggtgatcatc aactcctgcc tctgggatct ctcagatat ggtcgtgct caatggagag 420
ctaccgggag aacctggagc ggggtgttgt gcgcatggac caagtattgc cagactcctg 480
cctgctgggtg tggaaacatgg cgatgcccct cggggaacgt atcaactggg gtttcctcct 540
gccagagctc cagcccttgg caggctccct gcggcgggat gtggttgaag ggaacttcta 600
cagtgtctacg ctggccgggg accactgctt tgatgtccta gacctccact ttcacttccg 660
gcctgcagta cagcaccgtc atcgggatgg tgtccactgg gaccagcatg cacaccgcca 720
cctctcacac ctgcttctga cccatgtggc tgacgcctgg ggcgtggagc tgcccaagcg 780
tggctatccc cctgaccctg ggattgagga ctgggcagag atgaatcac cattccaggg 840
aagccatagg cagacccag acttcgggga gcacctggcc ttgctccac cccaccttc 900
ttcttttgct cctcccatgc cttttcccta cccgcttcc cagccctcgc cactccctc 960
cttcccccac ctgccccagg atacccttt tttccaggc cagcccttc caccctatga 1020
attcttcaac tataatccag tggaggactt ctgatgccca cccacttag gatgtggccc 1080
tggagtgaac tttgtgctg gccctctgcc acctccaac cctggcccta atcccatgg 1140
tcagcactgg ggcccagtg tccaccggg gatgccacgc tatgttcta acagccccta 1200
ccatgtgcgg agaattgggg ggccttgacg gcagcggctc agacactcag agagactgat 1260
ccacacatac aaactggaca gacggcctcc tgcccatcg gggacatggc ctgggtagac 1320
tggatcttgg gctgggactg gatgtgcaa tggccctca gggcctgcct ggcacctcag 1380
gtactgggct aggggtgtct ctatgcctgg tattgttct gtccattgct gtcaccaata 1440
aaggcatgga agaacagagt gaaaaaaaaa a 1471

```

&lt;210&gt; 25

&lt;211&gt; 3293

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1674661CB1

&lt;400&gt; 25

```

gcgagcgcgg gcaggcggcg acgcgggggc aggggtggac ggcggtcaga gccgaacgcg 60
agggcgggcg cgggggactg gagctgcgcg caataggaca gctggcctga agctcagagc 120
cggggcgctgc gccatggccc cactctgggc tgtctggctg ctggcagcaa ggctgtgggg 180
cctgggcatt ggggctgagg tgtggtggaa ccttgtgccc cgtaagacag tgtcttcttg 240
ggagctggcc acggtagtac ggcggttcta ccagaccggc atccaggact tctgacact 300
gacgctgacg gagcccactg ggcttctgta cgtggcgccc cgagaggccc tgttgcctt 360
cagcatggag gccctggagc tgcaaggagc gatctcctgg gaggcccccg tggagaagaa 420
gactgagtgt atccagaaaag ggaagaacaa ccagaccgag tgcttcaact tcatccgctt 480
cctgcagccc tacaatgctt cccacctgta cgtctgtggc acctacgctt tccagcccaa 540
gtgcacctac gtcacatgc tcaccttcac tttggagcat ggagagtttg aagatgggaa 600
gggcaagtgt ccctatgacc cagctaaggg ccatgctggc cttcttgggg atggtgagct 660
gtactcggcc aactcaaca acttctggg cacggaaccc attatcctgc gtaacatggg 720
gccccaccac tccatgaaga cagagtacct ggcccttttg ctcaacgaac ctactttgt 780
aggtctgccc tatgtacct agactgtggc cagcttcacg ggggacgacg acaaggctta 840
cttcttcttc agggagcggg cactggatgc cgtggcctat gccgagcagg tgggtgctcg 900
tgtggcccg gtctgcaagg gcgatatggg gggcgacagg accctgcaga ggaagtggac 960
cacgttcttg aaggcgcggc tggcatgctc tgccccgaac tggcagctct acttcaacca 1020
gctgcaggcg atgcacccc tgcaggacac ctectggcac aacaccacct tctttgggg 1080
ttttcaagca cagtggggtg acatgtacct ctggccatc tgtgagtacc agttggaaga 1140
gatccagcgg gtgtttgagg gccctataaa ggagtaccat gaggaagccc agaagtggga 1200
ccgtacact gacctgtac ccagccctcg gcctggctcg tgcattaaca actggcatcg 1260
gcgccacggc tacaccagct ccctggagct acccgacaac atcctcaact tctgaagaa 1320
gcaccccgat atggaggagc aggtggggcc tcgggtggagc cgccccctgc tctgaagaa 1380
gggcaccaac ttcaccacc tgggtggcga cggggttaca ggacttgatg gagccaccta 1440
tacagtgtct ttcattggca caggagacgg ctggctgctc aaggctgtga gcctggggcc 1500
ctgggttcac ctgattgagg agctgcagct gtttgaccag gagcccatga gaagcctggt 1560
gctatctcag agcaagaagc tgcctcttgc cggctcccg gtcctcgcct ggcagctgcc 1620
cgtggccgac tgcataagt atcgctcctg tgcagactgt gtcctcgcct gggaccctta 1680
ttgcgcctgg agcgtcaaca ccagccgctg tgtggcctg ggtggccact ctggatctct 1740
actgatccag catgtgatga cctcggacac ttcaggcatc tgcaacctcc gtggcagtaa 1800
gaaagtccag cccactccca aaaacatcac ggtggtggcg ggcacagacc tgggtgctgcc 1860
ctgccacctc tctccaact tggccccatg ccgctggacc tttggggggc gggacctgcc 1920
tgccgaacag cccgggtcct tctctacga tgcccggctc caggccctgg ttgtgatggc 1980
tgcccgagcc cgccatgcgg ggccctacca ctgcttttca gaggagcagg gggcgctgtg 2040
ggctgctgaa ggctaccttg tggctgtcgt ggcaggcccg tcggtgacct tggaggcccg 2100
ggcccccttg gaaaacctgg ggctggtgtg gctggcggtg gtggccctgg gggctgtgtg 2160
cctggtgctg ctgctgctgg tgctgtcatt gcgcggcgcg ctgcgggaag agctggagaa 2220

```

agggggccaag	gctactgaga	ggaccttggt	gtacccccctg	gagctgceca	aggagccccc	2280
cagtcceccc	ttccggccct	gtcctgaacc	agatgagaaa	ctttgggata	ctgtcggtta	2340
ctactattca	gatggctccc	ttaagatagt	acctgggcat	gcccgggtgc	agcccgggtg	2400
ggggccccct	tcgccacctc	caggcatccc	aggccagcct	ctgccttctc	caactcgggt	2460
tcacctgggg	ggtgggcgga	actcaaattg	caatggttac	gtgcgcttac	aactaggagg	2520
ggaggaccgg	ggagggctcg	ggcacccect	gcctgagctc	gcggatgaac	tgagacgcaa	2580
actgcagcaa	cgccagccac	tgcccgaact	caaccccag	gagtcatac	tatgagggga	2640
acccccaccg	cgtcggcggg	aagcgtggga	ggtgtagctc	ctacttttgc	acaggcacca	2700
gctacctcag	ggacatggca	cgggcacctg	ctctgtctgg	gacagatact	gcccagcacc	2760
caccgggcca	tgaggacctg	ctctgtctag	cacgggcact	gccacttggt	gtgggtcacc	2820
agggcaccag	cctcgcagaa	ggcatcttcc	tcctctctgt	gaatcacaga	cagcggggac	2880
cccagccgcc	aaaacttttc	aaggcagaag	tttcaagatg	tgtgtttgtc	tgtatttgca	2940
catgtgtttg	tgtgtgtgtg	tatgtgtgtg	tgcacgcgcg	tgcgcgcttg	tggcatagcc	3000
ttcctgtttc	tgtcaagtct	tcctttggcc	tgggtccctc	tggtagtca	ttggagctat	3060
gaaggggaag	gggtcgtatc	actttgtctc	tcctaccccc	actgccccga	gtgtcgggca	3120
gcgatgtaca	tatggagggtg	gggtggacag	ggtgctgtgc	cccttcaaaa	ggaagtgcaa	3180
ggccttgggg	tgggcctagt	cttgctcctt	agggctgtga	atggttttca	gaggttgggg	3240
gaaggaaatt	ggaaccctcc	tgtgtgtttt	gggggggaaa	gggttggttg	ggg	3293

&lt;210&gt; 26

&lt;211&gt; 1324

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1689337CB1

&lt;400&gt; 26

cagaatatca	gatgtccaac	atttcccttc	tggctctttt	tagctcccaa	gtatctgctt	60
ccttaaaggc	cctgagtcac	ttcttctctc	tttgcttcag	actggcccgg	gagcaagcgc	120
gagtgtgcga	actgcagagt	gggaaccagc	agctggagga	gcagcgggtg	gagctggttg	180
aaagactgca	ggccatgctg	caggcccaact	gggatgagge	caaccagctg	ctcagcacca	240
ctctcccgcg	gcccaccctt	ccagctcctc	ctgctggacc	ctccagcccc	gggcctcagg	300
agcccagaaa	ggaggagagg	agggctctga	ctatgcctcc	catggccgtg	gccctgaagc	360
ctgtattgca	gcagagccgg	gaagcaaggg	acgagctacc	tggagcgctt	cctgttcttt	420
gcagttcctc	ctcagatctt	agcctcctgt	tgggcccctc	ttttcagagc	cagcattctt	480
tcagccccct	ggagcccaaa	ccagacctca	cttcatccac	agctggggcc	ttctctgcac	540
ttggggcctt	ccatcccgat	catagggcag	aaaggccatt	ccctgaggaa	gacccctggac	600
ctgacgggga	gggcctccta	aagcaaggga	tgccgcctgc	tcagctggag	ggcctcaaga	660
atttttttga	ccagttgctg	gagacagtgc	cccagaacaa	tgagaaccct	tctgtcgacc	720
tgttgccccc	taagtctggg	cctctgactg	tcctcatctg	ggaggaagcc	cctcaagtgc	780
cacgtattcc	accgcctgtc	cacaaaacca	aagtccctt	agccatggca	tccagtcttt	840
tcgggttccc	tgagcctccc	tcctcccatt	cacaaggcag	tgggtcccag	agtggttccc	900
cagagagagg	tggagatggg	cttacattcc	caaggcagct	gatggagggtg	tctcaactgt	960
tgcgactcta	ccaggtctcg	ggctgggggg	ctctgcctgc	tgaggatctc	ctgtctctacc	1020
tgaagaggct	ggaacacagc	ggcaggactg	atggccgagg	ggataatgtc	cccagaagga	1080
acacagactc	ccgcttgggt	gagatcccc	ggaaagagat	tccttcccag	gctgtccctc	1140
gccgccttgc	tacagcccc	aagactgaaa	aacctcccgc	acggaagaaa	agtgggcacc	1200
ctgccccgag	tagcatgagg	agccgggggg	gagtctggag	atgagcccc	ctaccctctc	1260
tcctctttgt	tctctcattg	ttgttatttt	aataaatgct	cagtagtctg	taaaaaaaaa	1320
aaaa						1324

&lt;210&gt; 27

&lt;211&gt; 1884

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1746392CB1

&lt;400&gt; 27

cgaggggggtg	tggaaactta	ccggctgagc	catggataca	ccgttaagge	gcagccgagc	60
gctgggaggc	ctaaggcccg	aatcccccg	gagcctcacc	tcagtttcgc	ggacgagagc	120
ggcccttgtg	gagttcgagt	cgaaccacga	agaaacgagg	gagcccgggt	ctcctccgag	180
tgtgcagcgg	gctggcctgg	ggtcccccg	aaggccgcgc	aagacaagcc	caggatcacc	240

ccgtctgcag	caggggtgcag	gcttggagtc	accccaaggg	cagccagagc	caggcgcagc	300
gtccccccag	cgtcagcaag	acctacacct	ggagtcgcct	caaagacagc	cagagtacag	360
tcttgaatcc	ccacgatgtc	agccgaagcc	aagtggaggag	gcaccgaagt	gttctcagga	420
ccagggagta	ctggcctcgg	agtgggccc	gaataaggag	gagctgaccc	cgggggcccc	480
ccagcatcag	ctaccgccgg	tcccaggatc	accagagcct	taccccggtc	agcaagctcc	540
cggctccggag	ccctctcagc	cactactgga	gctgacaccc	agggcacctg	gtcccccccg	600
gggtcagcat	gagccagca	agccacctcc	agctggggag	acggtgacag	gcggcttcgg	660
ggcaaagaag	cgaaaaggtt	cttcatccca	ggcccagcg	tccaagaagt	tgaataaaga	720
ggagcttctt	gtaatcccg	aggggaagcc	caaatcgggg	cgagtgtgga	aggaccgctc	780
caagaaaaga	ttctcccaga	tgcttcagga	caagcccctg	cgcacatcgt	ggcagcggaa	840
gatgaaggaa	cgacaggaga	ggaagctggc	caaggacttt	gcccgtcacc	tggaggaggga	900
gaaggagagg	cgccgccagg	agaagaaaca	gcgcgggct	gagaacctga	aacgccgcct	960
ggagaatgag	cgggaaggcag	aggctcgtcca	agtgatccga	aaccccgcga	agctcaagcg	1020
ggcaaagaag	aagcagctgc	gctccattga	gaagcgggac	accctggccc	tgctgcagaa	1080
gcagccgccc	cagcagccgg	cagccaagat	ctgagctcag	gacggcccga	gggccttcca	1140
tggccaacaa	acatgtcaga	cacagcacct	caggccgctg	ctcagatgcc	tctgtctggag	1200
ctggcactcc	aaacccatgg	ctccagaaca	gggaccccca	ccccgaccgg	ggctcctcgg	1260
cctttgaagg	cttccaggca	ggtctgtgtg	ggacagaagc	ccagaggggg	cctgggacct	1320
ggcagagatg	ggggcgggaa	gagattcagc	tcccatecct	ccttctcttc	cttctccaag	1380
tgctttcaaa	ccaagaactg	tacattcttc	tggttccctc	gtgagctggg	gactggcagg	1440
tgactccctc	agcagtgtat	gccctttctc	agcctccctg	gtccatccca	ggcctggagg	1500
ctgacagtgg	ggaaatccagc	ttccccaca	ccttcccaaa	ggctgctctg	agcacctcca	1560
cacccactg	cctctgtccc	cagcaaactg	aatccggttc	ctctccactt	ttcaatactg	1620
aaagattaaa	atggggaggt	tgcagggagc	agagcttttc	cctagcacc	actttcccaa	1680
accagtctct	gcagaagccc	cagagaatct	aactcatgcc	tgctccagtct	acagcaaaaa	1740
tatttattga	gtgctgtgtg	catacaggca	caatcctagg	caccggcaaa	tacagacaat	1800
agaccaagtc	ccctgccctc	gaggagcttt	cattctgatg	gagagaaaac	ataataaaca	1860
agcaaaatgc	acaaaaaaaa	aaaa				1884

&lt;210&gt; 28

&lt;211&gt; 2429

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1825182CB1

&lt;400&gt; 28

cctattttcca	aaaagctcga	ctggagtggt	ataaaacctg	aaaattctct	tgtgctttct	60
cttctttttgc	ttctagttac	catcctcaaa	ggattggcta	aaagcaagca	actggattga	120
acaccctaag	aagaaagatt	cacactgcac	caggagacat	cagaaagaat	gaaaactctg	180
ccgctgtttg	tgtgcatctg	tgcactgagt	gcttgcttct	cgttcagtga	aggtcgagaa	240
agggatcatg	aactacgtca	cagaaggcat	catccaat	cacccaatc	tcactttgaa	300
ttaccacatt	atcctggact	gctagctcac	cagaagccgt	tcattagaaa	gtcctataaa	360
tgtctgcaca	aacgctgtag	gcctaagctt	ccaccttcac	ctaataagcc	ccccaaattc	420
ccaaatcctc	accagccacc	taaacatcca	gataaaaaata	gcagtgtggg	caaccctacc	480
ttagtggcta	caaccctaat	tccatctgtg	actttcccat	cagcttccac	caaaattact	540
acccttccaa	atgtgacttt	tcttcccag	aatgccacca	ccatatcttc	aagagaaaaa	600
gttaacacaa	gctcttctgt	agctacatta	gcaccagtga	attccccagc	tccacaagac	660
accacagctg	ccccaccac	accttctgca	actacaccag	ctccaccatc	ttctcagct	720
ccaccagaga	ccacagctgc	cccaccacac	ccttctgcaa	ctacacaagc	tccaccatct	780
tcttcagctc	caccagagac	cacagctgcc	ccaccacac	ctcctgcaac	taccagagct	840
ccaccatctt	cctcagctcc	accagagacc	acagctggcc	caccacacc	ttctgcaact	900
acaccagctc	cactatcttc	ctcagctcca	ccagagacca	cagctgtccc	accacacact	960
tctgcaacta	ccctagaccc	atcatccgcc	tcagctccac	cagagaccac	agctgcccc	1020
cccacacctt	ctgcaactac	accagctcca	ccgtcttccc	cagctccaca	agagaccaca	1080
gctgccccaa	ttaccacacc	taattcttcc	ccaactactc	ttgcacctga	cacttctgaa	1140
acttcagctg	caccacacac	ccagactact	acttcggtea	ctactcaaac	tactactact	1200
aaacaaccaa	cttcagctcc	tggccaaaat	aaaatttctc	gatttctttt	atatatgaag	1260
aatctactaa	acagaattat	tgacgacatg	gtggagcaat	agtatatgtt	atgttgtaaa	1320
gtgttctgtc	atttacaaga	tgtgattcat	gagtgcagaa	ctaccacctt	tcttttagca	1380
ccaatcccaa	catgaaatta	tattactcag	attttaaagc	ctatcattaa	tctttcaatc	1440
taattattca	ccaccacaag	acctattaac	aagacaaaat	gcctctatcc	cacaagccag	1500
atgcaggctc	ggggttcaaa	ataactcttt	ggatcctaca	gagatagcct	actgagggca	1560
gagaaagtc	ttagataaag	agagaatatt	gtatggggca	tcaaccattt	acttttccct	1620
gaatgttaga	aactacaaaa	ccactacctt	gtacccccat	caaaatccca	cctgaaccat	1680

ctaatectat	aaacataaag	gggtaaaatt	ggaactctcc	agatgaacaa	agacatctaa	1740
atatctgtag	atagaaacat	ttatctatct	aaatatattg	atagacctgt	cattgtattg	1800
attaatgaca	aaacccttta	gataattatc	ttccatttta	aataaaaattt	tatttcacaa	1860
atatgagcca	agaaagagga	aagttgattt	gaagtgagga	ttagaagtga	atgacaataa	1920
agtctggcag	ccaagcacga	accaagactg	gcactatttt	tcttagtgta	tataattggt	1980
taaaactgcaa	ggttgacatt	tattgtgttg	tgctaaagtt	aatttttgatc	taatgtacct	2040
gattctagcc	tctgtgaaca	acaagaatat	gtttgtgtat	gttcacatgg	tgcttataat	2100
atttcactat	caattcaatt	aattcacata	aattccatgt	gaaatgtatt	caacaatgga	2160
atattttcta	aaacatttag	tatacatttg	aatgtatttt	aaaccatgcc	aaactactgc	2220
tttaattgca	agtttgacga	attgtctctg	aaaataaaaa	ccctgacttt	agttgtaaaa	2280
caataaaaagt	tagctacttg	gtatacggag	atgttaattt	gggatatgga	ggcattttta	2340
tcttctgtca	ctactactta	aaactctgat	gattatgtta	gatttttttg	ctaactaata	2400
aagatttcaa	atggcaattt	ataaaaaaa				2429

<210> 29  
 <211> 985  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2155541CB1

<400> 29						
ggaagagggc	ggcgacggtg	gtggtgactg	agcggagccc	ggtgacagga	tggtgggtgtt	60
ggtattagga	gatctgcaca	tcccacaccg	gtgcaacagt	ttgccagcta	aattcaaaaa	120
actcctgggtg	ccaggaaaaa	ttcagcacat	tctctgcaca	ggaaaccttt	gcaccaaaga	180
gagttatgac	tatctcaaga	ctctggctgg	tgatgttcat	attgtgagag	gagacttcga	240
tgagaatctg	aattatccag	aacagaaagt	tgtagctgtt	ggacagttca	aaattgggtct	300
gatccatgga	catcaagtta	ttccatgggg	agatatggcc	agcttagccc	tggtgcagag	360
gcaatttgat	gtggacattc	ttatctcggg	acacacacac	aaatttgaag	catttgagca	420
tgaaaataaa	ttctacatta	atccaggttc	tgccactggg	gcatataatg	ccttggaac	480
aaacattatt	ccatcatttg	tggtgatgga	tatccaggct	tctacagtgg	tcacctatgt	540
gtatcagcta	attggagatg	atgtgaaagt	agaacgaatc	gaatacaaaa	aaccttaag	600
ccaggcctgt	cttgatgatt	tttggttttt	tttcatgttc	ctggtgaaat	caagtaatta	660
aacattttaag	agccacaaaa	ttgtatcact	tttataatat	tttgcagtaa	aatataatac	720
catcttctct	gttaatacat	aattgctcca	agcttctctg	aaactataag	aatatattta	780
gtttacagta	tatggattct	atgaaaaaat	gtccacaaca	cagtaattgg	tcacttgta	840
agaaaaaattt	atccttgtaa	gtatcttcaa	agttgatatt	tggaacttta	ttccaaaagt	900
agtgcatgtg	gagaaagaat	ctagactttc	ttgtatacat	ttttctcttc	tccagtaata	960
aacaattacc	tttcatttaa	aaaaa				985

<210> 30  
 <211> 3381  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2215706CB1

<400> 30						
gttagctcag	tgtaacattg	atgagctgaa	gaaagagatg	aatatgaatt	ttggagactg	60
gcaccttttc	agaagcacag	tactagaaat	gagaaacgca	gaaagccacg	tggtccctga	120
agaccacagt	ttcctcagtg	agagcagcag	tggtccagcc	ccgcacggtg	agcctgctcg	180
ccgcgcttcc	cacaacgagc	tgctcacac	cgagctctcc	agccagacgc	cctacacact	240
caacttcagc	ttcgaagagc	tgaacacgct	tggtctggat	gaaggtgcc	ctcgtcacag	300
taatctaagt	tggcagtcac	aaactcgcag	aaccccaagt	ctttcgagtc	tcaattccca	360
ggattccagt	attgaaattt	caaagcttac	tgataagggtg	caggccgagt	atagagatgc	420
ctatagagaa	tacattgctc	agatgtccca	gttagaaggg	ggccccgggt	ctacaacat	480
tagtggcaga	tcttctccac	atagcacata	ttacatgggt	cagagttcat	cagggggctc	540
tattcattca	aacctagagc	aagaaaaggg	gaaggatagt	gaaccaagc	ccgatgtgg	600
gaggaagtcc	tttctaata	agaggggaga	tgttatcgat	tattcatcat	caggggttcc	660
caccaacgat	gcttcccccc	tggtatcctat	cactgaagaa	gatgaaaaat	cagatcagtc	720
aggcagtaag	cttctcccag	gcaagaaatc	ttccgaaagg	tcaagcctct	tccagacaga	780
tttgaagctt	aagggaagtg	ggctgcgcta	tcaaaaactc	ccaagtgcag	aggatgaatc	840
tggcacagaa	gaatcagata	acactccact	gctcaaagat	gacaaagaca	gaaaagccga	900



```

agggaaagta gagagagtgc cgaagtctcc agaacacagt gctgagccga tcagaacctt 960
cattaaagcc aaagagtatt tatcggatgc gctccttgac aaaaaggatt catcggattc 1020
aggagtgaga tccagtgaag gttctcccaa tcaactctctg cacaatgaag tggcggatga 1080
ctcccatctt gaaaaggcaa atctcataga gctggaagat gacagtcaca gcggaaagcg 1140
gggaatccca catagcctga gtggcctgca agatccaatt atagctcggg tgtccatttg 1200
ttcagaagac aagaaaagcc cttccgaatg cagcttgata gccagcagcc ctgaagaaaa 1260
ctggcctgca tggcagaaag cctacaacct gaaccgaact cccagcaccg tgactctgaa 1320
caacaatagt gctccagcca acagagccaa tcaaaatttc gatgagatgg aggggaattag 1380
ggagacttct caagtcattt tgaggcctag ttccagtccc aacccaacca ctattcagaa 1440
tgagaatcta aaaagcatga cacataagcg aagccaacgt tcaagttaca caaggctctc 1500
caaagatcct ccggagctcc atgcagcagc ctcttctgag agcacagggt ttggagaaga 1560
aagagaaagc attctttgag aaaaacaagc aaaggagaag agtggtactg tacccttatg 1620
acagaattgt cctggatttt gactccatcc acgcccata cctttctaca ttttgctgac 1680
agataactaa ccgatgatga ggccgaggta aaagagacat ctgcagtgtg acagaaggga 1740
gcatgagaag catggctcac cagccagcct ctgtgtgctt tgtaattaga agcttcagaa 1800
ctcactaata ctactgtacc ttctattggc gcattacccc ataaaaacttt ttgagacgag 1860
gtgagatctg agtataaaga taggtcagaa gtatttttaa gggcttaatg tgccaaaaag 1920
aaaaaaagct agagaccctt tttgcaaaca tttgggtgacc acacatttga gggaagacgt 1980
ggcgttaggt gaagcagaag caaacctgc tcttaggggc tcacctaggt gagtgacag 2040
cctgtgacgc tacagggaga ggctgagtaa accgagatcc agcgttctgt atggcagggg 2100
tattgcttat cacagagggt ctgaagagta ggaagtacat aatgaagagg gctttaaaaa 2160
ttgccaaaca agtgagtcac cagggctggc agtagtgtga cggggctgtc ctgagctggt 2220
aggagagtat atgcggggag ggctgggtgac ctccgtgggt ttatatgtcg gaaactcttc 2280
tctccaaatc ccaggcctgg cttccagcac catccagctg tgcccaagaa gccacctgg 2340
tctgttctcc aactctttta aatggtgccc aacttttcta agtgagctta gcaatgagaa 2400
gaaaaaaa catgaattct ttttctggaa aatcaggggag acatgggtaa taataggtac 2460
taataaatat ttatagatga gtgaatgagg aaataattac atcaaaaagg tcagtgcaca 2520
ttgataaatg acaaggaaat atttaattag gtaaaactaa atcattgtct tctatactag 2580
gatagacttt atctacttca tctgttctca agtcagcatg ttagttctgg ggaaggatca 2640
taagaaagga aatacttttt aaaaaaaaaa ttggaaacat gtaacaaagc aagggtaaaa 2700
tatatatata tatctatata agtgctgtga ctgtaaaagt gtactttcca ttaattatta 2760
gccgagttaa gagaatggtc acattgaagt actgtgtgga ctagaaatgt accctgtcat 2820
catgcaatga aatattgtta tctgttttaac atagctcatt tatgtagaat gaattctggt 2880
ggtttacccc aagtcacagt taggacggta gatggtgaga tgcagatgc gctattatct 2940
agattcagtg ttacattttc gatgtttatc actcagtggg tttttattaa tatgctgatt 3000
aagttattta ctgggcagtg cattgtgcta aatagttgct cttttgtgtt tcattgcctt 3060
gatgtttgag tgtaatctag catttttaata cagtgtttat tttgcatgat ctttaacaaa 3120
tgtttttaagc aatttttaaaa aggcaggatg ttattgacat tatacactga agtcttaaca 3180
ttttaacatt tatagtgttt atttgcaaaa ttgtataatt aggaattatt tcagagacaa 3240
tgttttcttt ttcagggtgag tagttgccgc gtaatatcat tggagtacat tctttatact 3300
gtttgtgaaa ttaatactag catattaagt gtacaaatag atttagaaaa caataaaaaa 3360
ttgcatgcta aaaaaaaaaa a 3381

```

<210> 31  
 <211> 1803  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2347692CB1

```

<400> 31
gggctgttcc cggggagggt gtgatgggtt gacaggtgcg tgacagtggg agctgctctc 60
ggcacaagca tgtacggcaa aggcaagagt aacagcagcg ccgtcccgtc cgacagccag 120
gcccgggaga agttagcact ctacgtatat gaatatctgc tccatgtagg agctcagaaa 180
tcagctcaaa cattttttatc agagataaga tgggaaaaaa acatcacatt gggggaacca 240
ccaggattct tacattcttg gtggtgtgta ttttgggac tctactgtgc agctccagag 300
agacgtgaaa catgtgaaca ctcaagtga gcaaaagcct tccatgatta cagtgtgca 360
gcagctccca gtccagtgtc aggaacatt cccccaggag atggcatgcc agtagtctct 420
gtaccaccag ggttctttca gccttttatg tcacctcggt accctggagg tccaaggccc 480
ccattgagga tacctaatac ggcacttgga ggtgtcccag gaagtcagcc attactcccc 540
agtggaaatg atccaactcg acaacaagga catccaaata tgggtggggc aatgcagaga 600
atgactctc caagaggaat ggtgccctta ggaccacaga actatggagg tgcaatgaga 660
ccccactga atgcttttagg tggccctgga atgcctgga tgaacatggg tccaggtggt 720
ggtagacctt ggccaaaccc acaaatgcc aattcaatac catactctc agcatctcct 780
gggaattatg taggtcctcc aggaggtgga gggccaccag gaacacccat catgcctagt 840

```

```
ccagcagatt caaccaactc tgggtgataac atgtatactt taatgaatgc agtacctcct 900
ggacctaaaca gacctaatTT tccaatgggt cctgggtcag atgggtcccat ggggtggatta 960
ggaggaatgg agtcacatca catgaatggc tcttttaggct caggagatat ggacagtatt 1020
tccaagaatt ctcccaataa tatgagcctg agtaatcaac cgggcactcc aagggatgat 1080
ggcgaaatgg ggggaaatTT cttaaactct tttcagagtg agagttactc ccctagcatg 1140
acaatgagcg tgtgatccat taccaagctc cctcatgaaa accacagtga gtcagccctt 1200
cacagaacta ctacggaaga aaattattca tcacagtgtg cagttaaaca aaggaatctc 1260
agtcacacca aaccaacctt tttatttctt gctctctccc ctcttttgtg aagacagcg 1320
gtccaaatgt gattcaaaca actgtacgga gtggcatatt agaattgccc taaactgaac 1380
tgcaataaat tatgtgtgta tgtatatgtg tgggaaagag aatgtactgt atatgtgtat 1440
gtttatcaga catatacaca tacatacatt gacccacagg acattgtaaa atattatcac 1500
atgacatctt aagtagaaat aagtagggac ttttattcca tccttttttt cactgtttaca 1560
ttttaattat tacaagttgc tcctgcccc tcctgaact attttgtgct gtgtatatca 1620
ctgctttata taagttatTT ttttaagggtga actcagatgt tatggttttg taaatgtctg 1680
caatcatgga taggaataaa atcgcttatt tgagagcttt cattaaattg cgtctgatgc 1740
aagttatcct gtgaatcaca aagtgtactg tctcaaaaaa tagaagaaaa tgtgaaaaaa 1800
aaa 1803
```

<210> 32  
<211> 1515  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 2579048CB1

```
<400> 32
ggtaaagtgg cttctgggcg gaaggtacac tataggctcg gggaggtaag cggcggcagg 60
cggcggttg gtgtgtccc ggtgtgggga ggcgacagag ccctggcact tgagggttga 120
gggggcctcc ccagcgcggc gaaccgtcta gctcccgag gccaggccgt gactgcggga 180
ggtatagccc aaggcggaag aattttgcca ctactacct gtgtgacctc gggtaaatta 240
gccttggaac gtcagtttct tcgtctctat aattgaaata ataatagtag ctctctcagg 300
attgtttgta gccgtcagtg aaacacttag agcagtttct ggcacatggt agaattgggc 360
tatttgcctga agcttcttgg tggcccttgc tagcccagga agaaacttac attttgattt 420
ttttgtacca tggcttttgg tcacaaattg ctgcatggta cttattttct cagaaaattc 480
tctaagccaa cttctgcctt gtatccattt ttgggtattc tctttgcaga gtattccagt 540
agtcttcaga aaccagtggc ttctcctggc aaagcctcct cacagaggaa gactgaaggg 600
gatttgcaag gatcacca gaaagaagtt gctttggata taacttcttc tgaggagaag 660
cctgatgtta gtttcgataa agcaattaga gatgaagcaa tataccattt taggcttttg 720
aaggatgaaa ttgtggatca ttggagagga ccggaaggcc accctctgca tgagggtctg 780
ctggaacaag ccaaggttgt ctggcaattc cgggggaaag aagatttgga taagtggaca 840
gtgacttctg ataagacgat tggaggcaga agtgaagtgt ttttgaaaat gggcaagaat 900
aaccaaagtg cactgctata tggaaactcg agctctgagg cgctcagga cggggagtct 960
acccgaagtg ggtactgtgc aatgatattc aggtatccaa ggggtgcttt tgagaggaag 1020
atgtcttacg attggtccca gttcaatact ctgtatctcc gtgtacgtgg ggatggtcgg 1080
ccttggtatg tgaatatcaa ggaggacaca gatttcttcc agaggacgaa tcagatgtat 1140
agttacttca tgttcacccg cgggggaccc tactggcagg aggtcaagat tcctttttcc 1200
aaatttttct tctctaatec aggaagaatc cgggatgttc agcatgagct tccgcttgat 1260
aagatctctt ctataggatt caccttggct gataaagtgg atggtccatt ctctctggag 1320
atagatttta ttggcgtgtt tactgatcca gtcatacag aagaatttgc ctatgaaaat 1380
tctccagagc ttaacccaag gcttttttaa taaagatcat atggtagtgt tgttttacta 1440
atctaagggt actagatctt acaatgatat agacaaaata aaatatttct ttaatggcat 1500
ccaaaaaaa aaaaa 1515
```

<210> 33  
<211> 4416  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 2604493CB1

```
<400> 33
cggatgggcg ggcagccgcg ccgcgcgggc acttttttaa ttttttcggg tgccgcagcg 60
gcgacccctc ggcgcgatg tcctgatcc ctggagcgac gacggccgct gcctaagctg 120
```

gaaagaggaa	tgccagctcc	tgagcaggcc	tcattggtgg	aggaggggca	accacagacc	180
cgccaggaag	ctgcctccac	tggcccaggc	atggaacccg	agaccacagc	caccactatt	240
ctagcatccg	tgaaggagca	ggagcttcag	tttcagcgac	tcacccgaga	actggaagtg	300
gaaaggcaga	ttgttgccag	tcagctagaa	agatgtaggc	ttggagcaga	atcaccaagt	360
atcgccagca	ccagctcaac	tgagaagtca	tttcccttga	gatcaacaga	cgtgccaaat	420
actggtgtaa	gcaaacctag	agttttctgac	gctgtccagc	ccaacaacta	tctcatcagg	480
acagagccag	aacaaggaac	cctctattca	ccagaacaga	catctctcca	tgaaagttag	540
ggatcattgg	gtaactcaag	aagttcaaca	caaatgaatt	cttattccga	cagtggatac	600
caggaagcag	ggagtttcca	caacagccag	aacgtgagca	aggcagacaa	cagacagcag	660
cattcattca	taggatcaac	taacaacccat	gtggtgagga	attcaagagc	tgaaggacaa	720
acactggttc	agccatcagt	agccaatcgg	gccatgagaa	gagttagtcc	agttccatct	780
agagcacagt	ctccttctta	tgttatcagc	acaggcgtgt	ctccttcaag	ggggtctctg	840
agaactttct	tgggtagtgg	atttggctct	ccgtcagtga	ccgacccccg	acctctgaac	900
cccagtgcac	attcctccac	cacattacct	gctgcacggg	cagcctctcc	gtactcacag	960
agacccgcct	ccccaacagc	tatacggcgg	attgggtcag	tcacctcccg	gcagacctcc	1020
aatcccaacg	gaccaacccc	tcaataccaa	accaccgcca	gagtgggggtc	cccactgacc	1080
ctgacggatg	cacagactcg	agtagcttcc	ccatcccaag	gccaggtggg	gtcgtcgtcc	1140
cccaaagcgt	cagggatgac	cgccgtacca	cagcatctgg	gaccttccct	gcaaaggact	1200
gttcatgaca	tggagcaatt	cggacagcag	cagtatgaca	tttatgagag	gatggttcca	1260
cccaggccag	acagcctgac	aggcttacgg	agttccctatg	ctagtcagca	tagtcagctt	1320
gggcaagacc	ttcgtttctgc	cgtgtctccc	gacttgcaca	ttactcctat	atatgagggg	1380
aggacctatt	acagcccagt	gtaccgcagc	ccaaacctag	gaactgtgga	gctccaagga	1440
tcgcagacgg	cgttgtatcg	cacaggttca	ggtattggaa	atctacaaag	gacatccagc	1500
caacgaagta	cccttacata	ccaaagaaat	aattatgctc	tgaacacaa	agctacctac	1560
gcggagccct	acaggccctat	acaataccga	gtgcaagagt	gcaattataa	caggcttcag	1620
catgcagtgc	cggtctgatga	tggcaccaca	agatccccat	caatagacag	cattcagaag	1680
gaccccgagg	agttttgcctg	gcgtgtacct	gagttgcctg	aggctattca	catgcttcag	1740
caccagttcc	catctgttca	ggcaaatgca	gcggcctacc	tgcagcacct	gtgctttggg	1800
gacaacaaag	tgaagatgga	ggtgtgtagg	ttagggggaa	tcaagcatct	ggttgacctt	1860
ctggaccaca	gagtttttga	agttcagaag	aatgcttgtg	gtgcccttcg	aaacctcggt	1920
tttggcaagt	ctacagatga	aaataaaaata	gcaatgaaga	atgttgggtg	gatacctgcc	1980
ttgttgcgac	tgtttgagaaa	atctattgat	gcagaagtaa	gggagcttgt	tacaggagct	2040
ctttggaatt	tatectcatg	tgatgctgta	aaaatgacaa	tcattcgaga	tgctctctca	2100
accttaacaa	acactgtgat	tgttccacat	tctggatgga	ataactcttc	ttttgatgat	2160
gatcataaaa	ttaaatttca	gacttcacta	gttctgcgta	acacgacagg	ttgcctaagg	2220
aacctcagaa	cgcgggggga	agaagctcgg	aagcaaatgc	ggtcctgcga	ggggttgtaa	2280
gactcactgt	tgtatgtgat	ccacacgtgt	gtgaacacat	ccgattacga	cagcaagacg	2340
gtggagaact	gcgtgtgcac	cctgaggaac	ctgtcctatc	ggctggagct	ggaggtgccc	2400
caggcccggt	tactgggact	gaacgaattg	gatgacttac	taggaaaaga	gtctcccagc	2460
aaagactctg	agccaagtgg	ctgggggaag	aagaagaaaa	agaaaaagag	gactccgcaa	2520
gaagatcaat	gggatggagt	tggtcctatc	ccaggactgt	cgaagtcccc	caaagggtgt	2580
gagatgctgt	ggcaccatc	ggtggtaaaa	ccatatctga	ctctcttagc	agaaagtccc	2640
aaccagccca	ccttggaagg	ctctgcaggg	tctctccaga	acctctctgc	tggcaactgg	2700
aagtttgcag	catatatccg	ggcggccgtc	cgaaaagaaa	aggggctccc	catccttgtg	2760
gagcttctga	gaatggataa	cgatagagtt	gtttcttccg	tggcaacagc	cttgaggaa	2820
atggcactag	atgttcgcaa	caaggagctc	ataggcaaat	acgccatgcg	agacctggct	2880
aaccggctcc	ccggcggaac	tggcccccag	gtcttgtctg	atgagaccat	ggcagccatc	2940
tgctgtgctc	tgacagaggt	caccagcaaa	aacatggaga	acgcaaaagc	cctggccgac	3000
tcaggaggca	tagagaagct	ggtgaacata	accaaaggca	ggggcgacag	atcacctctg	3060
aaagtgggtga	aggcagcagc	ccaggctctg	aatacatat	ggcaatatcg	ggacctccgg	3120
agcatttata	aaaaggatgg	gtggaatcag	aaccatttta	ttacacctgt	gtcgacattg	3180
gagcgagacc	gattcaaatc	acatccttcc	ttgtctacca	ccaaccaaca	gatgtcacc	3240
atcattcagt	cagtcggcag	cacctcttcc	tcaccagcac	tgtaggaat	cagagaccct	3300
cgctctgaat	acgataggac	ccagccacct	atgcagtatt	acaatagcca	aggggagccc	3360
acacataaag	gcctgtaccc	tggtaagacg	ccagttgggt	gtgtgatata	gtctcttgaa	3420
aagccgcatt	tccaggcgct	tggccagtg	cctgggaagt	agcctgtgct	tgtattgaga	3480
cagtcceccc	gcagcaaacc	atgttccagt	cattcccttt	cctactttgg	ggattgttgc	3540
cttttctgct	tgtttaaagt	aaaacaagca	tgtactttgt	tgtatgtatg	tatgtatgta	3600
gttgtacggt	gggcacaaat	aaaaagaggg	ctgtatccaa	ataaatcatt	tctggctgct	3660
cactggcaca	gtccctttgc	tccgtccctc	cctggctcga	gcagtcctctg	tgttttccac	3720
atgcatcaga	accgtcagcc	cagtgctcact	ttgcaggggc	cacatcttct	agactggctc	3780
atctcttaaa	ttcaaaccag	agaatggaga	tcaatttcat	ttacattttc	atggagaaga	3840
gtttgaataa	atagcaaaa	gtataatgtg	attcttctca	acgtgattat	tttttaagaa	3900
taacaacaaa	aaataatggt	acatagtgtt	aatattttta	tatcctattc	agaaagctgt	3960
gcatacaatt	tatttaaagg	ataccagccc	aaagggttac	taagatttca	tactgttccc	4020
attctcaacc	gtctggcgat	tataactact	ttagatgttc	caaataagtc	agtgtgagct	4080
accccaaaa	aagtactctt	ttctcaaacc	tgtctattaa	aaataaataa	aagctgggtg	4140

```

cagctttact cctgctgggg gggcagtcct cattgtactg ttgtctttga aattgtgatt 4200
tcacaggact atttactaat cgactgttgt tactatggca attaatagct cagaaaattg 4260
cagatgctac atttcaacac caagctctgg gcaaaaaggaa atgcacatac caactgaagt 4320
gcatgggttg ttcgattttg cagcagcctg ctttatctta cagatcacct ctttagcatg 4380
catgcccttc tgtatcttct tcgcttgctt caataa 4416

```

```

<210> 34
<211> 4428
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 2787182CB1

```

```

<400> 34
cggaggggccg gggctgggtg gcagcggcca tggaccgctt cgtttggacc agcggcctcc 60
tggagatcaa cgagaccctg gtgatccagc agcgcggggg gcgaatctac gatggcgagg 120
agaagataaaa atttgatgct gggactctcc ttcttagtac acaccgactg atttggagag 180
atcagaaaaaa tcatgagtggt tgcattggcca ttctctcttc ccaaattgtg ttcatggaag 240
aacaggcggc tggaattggg aagagtggca aaatagtggg tcattcttcac ccagctcctc 300
ctaacaaaga acctggccca ttccagagta gtaagaactc ctacatcaaa ctctccttca 360
aagaacatgg ccagattgag ttttacaggc gtttatcaga ggaaatgaca caaagaagat 420
gggagaatat gccagtttcc cagtcattac aaacaaatag aggaccccag ccaggaagaa 480
taagggctgt aggaattgta ggtattgaaa ggaaactgga agaaaaaaa aaagaaactg 540
acaaaaacat ttctgaggcc ttggaagacc tcagcaaact aatgatcaag gctaaggaaa 600
tgggtggaatt atcaaaatca attgctaata aaattaaaga caaacaagggt gacatcacag 660
aagatgagac catcagggtt aaatcctact tgctgagcat gggaatagct aaccagttta 720
ccagagaaac ctacggctca ggcacacagt accacatgca gctggccaaa caactggctg 780
gaatattgca ggtgccttta gaggaacgag ggggaataat gtcactcacg gaggtgtact 840
gcttagtaaaa ccgagctcga ggaatggaat tgctctcacc agaagattta gtgaatgcgt 900
gcaagatgct ggaagcactg aaattacctc tcaggtctcg tgtgtttgac agtggcgctca 960
tggttaattga gcttcagctc cacaaggaag aggaaatggt ggcctcggcc ctggagacag 1020
tttcagaaaa gggatcccta acatcagaag agtttgctaa gcttgtggga atgtctgtcc 1080
tcctagccaa agaaaggttg ctgcttgca gaaagatggg ccactcttgc cgtgatgact 1140
cagtgggaagg cctgcgtttt tacccaaatt ttttatgac acagagctaa ggtttttgta 1200
tttaaaatcc tttttgtcca tatgcttgcg tcattgtagag gttgtatgac attgagctaa 1260
gagataaacc ccgatcaatt gagaatttat tagaacttca cagtgcgaatg taaatctctt 1320
ttaatttctc actaaatag gtccaggaaa ttattttagt atacgcatag gaaaattcag 1380
aaaagtgaat gccaatatga atttaaaatc atgctatagt gcagaaccct cagagtttaa 1440
cttggaatat agtggatttt aacttgatcc tcaaatctaa tcattttata aagaagggaa 1500
tttagttttg cagagaataa aaagagaagt tgcatgttca gacagggttag attattattt 1560
tggtgtaact gaaattcact gattgcacat gacaatgttg ggacaaaata tactgcagca 1620
tgctatatga ggtctctccc cagggtttt agaagcagtc atagacatgt cttcaacata 1680
ccaaataaaa taccttttaa aatgaaataa ttttatttga cacatatatt tatatatatt 1740
ctatctagtt tctctttgtt tttttaagtg atgatttcat gactggcatt taaagaatgc 1800
aactgtgtca ttttgtttcc aaatgctgtg gattttgaaa ctgaactaga gagctgtata 1860
gacatgcccc gacttatgat tacaaattta ggaggtagac ggctcaggaa ttccctggga 1920
ttgttgtgct ggtggaatgg cagagggaac ttacacggaa ccttagtggt cttttacctc 1980
aaagccacag acaggaaata gaaagtggaa aagtaatatc tctttttctt ttccataagg 2040
agtttcaaca ctgaacttta aaaagtctat catattccag caatattttt tctttgtcct 2100
ttatgttgta agttgtgtgg aaaaactact tcggtaagaa atgttactga gataacaaca 2160
actggctaata actgcatgta gattgcttag gttttaaagt gactgcctga cttcacatgt 2220
tattgctaca gcctccagta tgcttcgcat atctcaaaact cagggacccc acaggacagg 2280
agacaccctt tctgaaactg agttggaagt gaaaggggtg tgatggtttt ggccaagcct 2340
gcgggaggga aagtattgta ttgggagcac ccttggggacc aggaagaggg atgccagggt 2400
tcacactctg ggacccttaa gacattgtca gtgggtaagg tggaggggcca ctgccaggag 2460
ctgctgcagt gccctctggg aagtgccagg gacctcagc tgcagacca ggggatccca 2520
atcccttate tctgctgaa aagtggctat tgctgctcg tctctccag tgcacagga 2580
atgctgtgcc tgagacatcc cactttagtt tttgttttcc aaacctgtca ctgactcttc 2640
attgtgggtga cagatcctct ggttccccca ttgctcagcc tcagggttat tttgagggtta 2700
actcagtgtc tgactgccc aagtcttttag gggttagttg gaccattgtg taagcttgtt 2760
tgtatgtctc acctctttct ggttggtact tactgtctca cactgctctg gcaaacttgt 2820
acacacacac acactctctc tctctctcac ctggctaagg cttttaaaat tatgaagata 2880
aataatctgt ttcaccagct ggagtgaagt gcaaggaaga ttgctggagc tactcaatct 2940
agcgctaata gtttggattc attactgcaa acctacataa tttacatat ttgtttactt 3000
tagtgtgaca actgatgaaa aaaaatggag caatctgaat tgtataaaat aacttaagaa 3060

```

```

ggaagaaaag tgatatataa atatattttt caaatgtcac attaatttaa aaatgagtat 3120
gattgatttt atttttaaag tgggcattct tcactgtttc gagacctttg tatgtatttg 3180
tgtattttta tctttttttt tcaggccatt ttataaagg gttatttttg cctctaatg 3240
tagaagttat gtttaataac aaccaatcag gccctaagtg caaaaaagtt ctgtcttggt 3300
gcttatcact ggtataatta ttattttttt gttttcctaa ctttgctctt aggagcatga 3360
gcctctttgt agctttatgg taatgcaata tttctggtea tttactgtca aaaaattttt 3420
ttacattgtg ttagtgaaga ctgtgttttc aggggttaa gattgggtatc aatgtatata 3480
gagtaaaatt attgcaaaat ttaaaaatga tttttcttgg cattcttttt aaatactctg 3540
aaacatatat gcaaatgaga aacctttaat catactaaag ccagttatgt taagaatttt 3600
tcctttggat ttataatta agagttgctt ataaatactt atgacattga gctcttactg 3660
ttttagttgt cttaaagaa tgggacatgt catgaactgg tttctttttt taaaaaattc 3720
ttgatgtgaa ttccactgtt acctgacct gtatattcag atgaatgtac tctttcttgt 3780
tccttggtac cagattaatc tgagagaatg tggattcacc ctgtcatctt tctcactac 3840
aaatacttaa aatggttctt ttgccccaaa gtctcagagt cttcattttg gtacagtggg 3900
gtcacaaact gtaagggtgt taggttttaa agaccgtgcc taaggacaaa attgagcatg 3960
ctcagaatgg tccttgaaaa tctcctaaat taccaataaa tttctctttg tagtcttaa 4020
gtttaaatcc ctgtggagtg atatgtaggt atgaatata tatgcttata ttaagtacaa 4080
aaatacaaac tgcataatca gagatttcta tagcattaat aatttccatg catgtgtctt 4140
ttccagtag gtatggttga atttatgtaa atttattgct aatcccatcc cttacgatct 4200
agagtataag ctgcgcaagg gcagaagttt ttatctggtt tgttcattga tgtattctaa 4260
gagctgagaa cagggcctgg acacaataag cattcaataa atatttactg aatgaatgaa 4320
ctcctaccta tattcctatt tataatttgg ctccacttta tctacttta gctcccatc 4380
aattcaataa aaaaaacatt tttttgaaca cataaaaaaa aaaaaaaa 4428

```

&lt;210&gt; 35

&lt;211&gt; 1907

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3096668CB1

&lt;400&gt; 35

```

gcgttgctgc gtaaatggcg ggggcgtgtc ttttggctcc tccgcgtgta gttacctgag 60
aaacgcggga agttgggccc aggcagtgtt gctgcggttg cctaagttgt tttactattt 120
ctggagagag ccgtgagctt gtccaggggc cccaatcctg aggcgcaccc ggtttctggc 180
gcggtgcgat ggaggtagtg gaggcgcg cgctcagct ggaaactctg aaattcaatg 240
gcaccgactt tggagttggg gaaggtccgg cggctccgtc tccgggctct gcccctgtgc 300
cagggacaca gccgcgcta cagtcgtttg aggggtcccc ggacgctggg cagaccgtgg 360
aggttaagcc tgccggggag cagcctctgc agcccttct gaacgcgctc gcggccgga 420
ccccggcgcc gcagccacag ccaccggctg aatcgccggc ctgcggagac tgcgtcacc 480
ccccaggagc cgcagagcct gcgcggggcg cggactcctt ggagacctcg gactcggatt 540
cggactcgga cagtgaaca gattcagata gttcaagttc atcgtcttcc tcttcatctt 600
cctcatcgcc gtcttcttcc tcttgtaata ctttctccc agtgcgtgca gatggagatg 660
atgatttaca agttgagaag gaaaataaga attttctct taaaaacaaa gatgaattac 720
ttcttaatct atagagaaaa aaaaaattca gtcattggct tcagtactcc agggattgga 780
tctactttc caaccatata tgaaaagtga aggtcacct aatagcatca ggtaattggc 840
tgccacgtgc atgtcctgtg ttgggtaata tgagaaaagt ggtagataa tttcatccca 900
ccttctcac ttttctcaat tctcccaaaa ctgacaccaa aaaggtttag ttttcatctg 960
gaatgtttga ttccttctcc caccaacctc tgttgctctt tttccatatt tgtgagtga 1020
aatcctactg aaagtctctt atataatttt actttagaaa gtcattcaatt tactgatact 1080
cttttttgc caataagcct tcacaaatgc ttgaaaagtc aacagtgtgg taaatgataa 1140
atgtttacct ttcagagggc cacttttgac atattagctt cttccatttt tgcattgaaa 1200
gtgttaggca cccatttttt gactggatg tctataacct catttttaat taggatggat 1260
tttttaatgt tatgtggtac tcttttgaat attttgctta actgaataac cttgtagacc 1320
atacagatgt tagaaaatta aatatataata atcttacaaa aacagacaaat aattgctctt 1380
tcattgatat tccagaagtg tggtaggat cttgcatttc tttaatgcca tcgttacgag 1440
tatctgttat ccagaagca cagtaatttt tttattggga tctagtatat aatttctgta 1500
ttaatgtgtt gtctttttga ttacatgcag ttggacattt aaattgtcgt tagccatttg 1560
tgtaactgtt tagaacttgt gtcagtttc cttactatc gtaggtaaga aaaatgaaac 1620
cagaaagaaa gagacttgt tgaagtcata ttagtgggga ggagcagatc tgaactggta 1680
atagatacat tccaactgc tgtgttttta gttattgtgg agacctgcat agcaacgttg 1740
gaatctggtc gtttggactt tctcattatt tggttctgta ataactatgt agcatatttt 1800
ctcaaagtga aggatgaatg tatcatctaa tgtaagcttt ttaaaattgt atttgcttgg 1860
attatattgc ctctcaataa agcttttgac ttggaaaaaa aaaaaaa 1907

```

<210> 36  
 <211> 1839  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3143411CB1

<400> 36  
 gccggctccgtc ggtggcctag agatgctgct gccgcgggtg cagttgtcgc gcacgcctct 60  
 gccgcgccagc ccgctccacc gccgtagcgc ccgagtgtcg gggggcgcac ccgagtcggg 120  
 ccatgaggcc gggaaaccgc ctacaggccg tgcgtctggc cgtgctgctg gtggggctgc 180  
 gggccgcgac gggctgcctg ctgagtgggc agccagtcctg ccggggaggg acacagaggc 240  
 cttgtttataa agtcattttac ttccatgata cttctcgaag actgaacttt gaggaagcca 300  
 aagaagcctg caggagggat ggaggccagc tagtcagcat cgagtctgaa gatgaacaga 360  
 aactgataga aaagtccatt gaaaacctct tgccatctga tggtgacttc tggattgggc 420  
 tcaggaggcg tgaggagaaa caaagcaata gcacagcctg ccaggacctt tatgcttgga 480  
 ctgatggcag catatcaciaa ttttaggaact ggtatgtgga tgagccgtcc tgcggcagcg 540  
 aggtctgcgt ggtcatgtac catcagccat cggcaccgcg tggcatcgga ggcccctaca 600  
 tgttccagtg gaatgatgac cgggtgaaca tgaagaacaa tttcatttgc aaatattctg 660  
 atgagaaacc agcagttcct tctagagaag ctgaagggtg ggaaacagag ctgacaacac 720  
 ctgtacttcc agaagaaaca caggaagaag atgccaaaaa aacatttaaa gaaagtagag 780  
 aagctgcctt gaatctggcc tacatcctaa tccccagcat tccccctctc ctctctcttg 840  
 tggtcaccac agttgtatgt tgggttttga tctgtagaaa aagaaaacgg gaggagccag 900  
 accctagcac aaagaagcaa cacaccatct ggccctctcc tcaccaggga aacagcccgg 960  
 acctagaggt ctacaatgtc ataagaaaac aaagcgaagc tgacttagct gagaccggc 1020  
 cagacctgaa gaatatttca ttccgagtgt gttcgggaga agccactccc gatgacatgt 1080  
 cttgtgacta tgacaacatg gctgtgaacc catcagaaag tgggtttgtg actctggtga 1140  
 gcggtggagag tggattttgt accaatgaca tttatgagtt ctccccagac caaatgggga 1200  
 ggagtaagga gtctggatgg gtggaaaatg aaatatatgg ttattaggac atataaaaaa 1260  
 ctgaaactga caacaatgga aaagaaatga taagcaaaat cctcttattt tctataagga 1320  
 aaatacacag aaggtctatg aacaagctta gatcaggtcc tgtggatgag catgtggtcc 1380  
 ccacgacctc ctgttggaac cccacgtttt ggctgtatcc tttatccag ccagtcaccc 1440  
 agctcgacct tatgagaagg taccttgccc aggtctggca catagtagag tctcaataaa 1500  
 tgtcacttgg ttggttgtat ctaactttta agggacagag ctttacctgg cagtataaaa 1560  
 gatgggctgt ggagcttgga aaaccacctc tgttttccct gctctataca gcagcacata 1620  
 ttatcataca gacagaaaat ccagaatctt ttcaaagccc acatatggta gcacaggctg 1680  
 gcctgtgcat cggcaattct catatctgtt tttttcaaag aataaaatca aataaagagc 1740  
 aggaaacaga aaaaaaaagg agatagggtg gtagatgatg gtaaggagtg ctaagaagg 1800  
 tgtgggaaga gttgattgaa tggagggggt cccttgttt 1839

<210> 37  
 <211> 503  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3170835CB1

<400> 37  
 cggctcgaga tgaaattcgc catcgtcctg ttgcacctct ttgccgtggc cctggctgcc 60  
 cctactgtcg aggtcctgcg atcggatagc aatggttgaa tcgataacta ctcatatgca 120  
 gttgaaacca gcgacggtag ctcgaagagc gaggagggtg tgctgaagaa cgccggcacc 180  
 gagctagagg ccattctcaac ccacggctcc ttcagctacg tgggccctga tggccagacc 240  
 tacaccgtca cctacgtggc cgatgagaac ggattccagc cccagggtgc tcatctgccc 300  
 gttgcccccg ttgcctaagc taaggagtgt aaataaattg tttataaaaag caaaaaaact 360  
 ggcagaaaaga acccaaagta cgtagacctt catgctagag agaattgtat gtctgccgaa 420  
 gacaaccgta ccgcctgcg caatcttacg gccactagtg gatcctgcgc atgaggacac 480  
 cgagagcgag cgtggggggg ggc 503

<210> 38  
 <211> 2154  
 <212> DNA  
 <213> Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3550808CB1

&lt;400&gt; 38

```

gaggacagga tgaggcccg cctctcattt ctctagccc ttctgttctt ccttggccaa 60
gctgcagggg atttggggga tgtgggacct ccaattccca gccccggctt cagctctttc 120
ccaggtgttg actccagctc cagcttcage tccagctcca ggtcgggctc cagctccagc 180
cgcagcttag gcagcggagg ttctgtgtcc cagttgtttt ccaatttcac cggctccgtg 240
gatgaccgtg ggacctgcca gtgctctgtt tccctgccag acaccacctt tcccgtggac 300
agagtgaac gcttgaattt cacagctcat gttctttctc agaagtttga gaaagaactt 360
tctaaagtga gggaatatgt ccaattaatt agtgtgtatg aaaagaaact gttaaactta 420
actgtccgaa ttgacatcat ggagaaggat accatttctt acactgaact ggacttcgag 480
ctgatcaagg tagaagtga ggagatggaa aaactgggtca tacagctgaa ggagagtttt 540
ggtggaagct cagaaattgt tgaccagctg gaggtggaga taagaaatat gactctcttg 600
gtagagaagc ttgagacact agacaaaaac aatgtccttg ccattcgccg agaaatcgtg 660
gctctgaaga ccaagctgaa agagtgtgag gcctctaaag atcaaaacac ccctgtcgtc 720
caccctcctc ccactccagg gagctgtggt catggtgggt tgggtgaacat cagcaaacgg 780
tctgtggttc agctcaactg gagagggttt tcttatctat atggtgcttg gggtagggat 840
tactctcccc agcatccaaa caaaggactg tattgggtgg cgccattgaa tacagatggg 900
agactgttgg agtattatag actgtacaac acactggatg atttgctatt gtatataaat 960
gctcgagagt tgcggatcac ctatggccaa ggtagtggta cagcagttaa caacaacaac 1020
atgtacgtca acatgtacaa caccgggaat attgccagag ttaacctgac caccaacacg 1080
attgctgtga ctcaaaactc ccctaattgt gcctataata accgcttttc atatgcta 1140
gttgcttggc aagatattga ctttgcgtgt gatgagaatg gattgtgggt tatttattca 1200
actgaagcca gcactggtaa catggtgatt agtaaaacta atgacaccac acttcagggt 1260
ctaaacactt ggtataccaa gcagtataaa ccatctgctt ctaacgcctt catggtatgt 1320
gggtttctgt atgccacccg tactatgaac accagaacag aagagatttt ttactattat 1380
gacacaaaca cagggaaga gggcaacta gacattgtaa tgcataagat gcaggaaaaa 1440
gtgcagagca ttaactataa cccttttgac cagaaacttt atgtctataa cgatgggtac 1500
cttctgaatt atgatctttc tgtcttgacg aagccccagt aagctgttta ggagttagggt 1560
tgaaagagaa aatggtttgt gaaaaaatag tcttctccac ttacttagat atctgcagggt 1620
gtgtctaaaa gtgtgttcat ttgagcaga tgtttagggt catagtctta ccacactaga 1680
gatctaggac atttgtcttg atttgggtgag ttctcttggg aatcatctgc ctcttcaggc 1740
gcattttgca ataaagtctg tctaggggtg gatgttcaga ggtctagggg cactgtgggg 1800
ctagtgaagc ctactgtgag gaggttcagc tagaagcctt aaattaggaa ttaaggaaact 1860
taaaactcag tatggcgtct agggattctt tgtacaggaa atattgcccc atgactagtc 1920
cccatccatg tagcaccact aattcttcca tgctggaag aaacctgggg acttaggttag 1980
gtagattaat atctggagct cctcgaggga ccaaactctc aacttttttt tccccctact 2040
agcacctgga atgatgctt gtatgtggca gataagtaaa tttggcatgc ttatatattc 2100
tacatcttga aagtggcctg gttttatgga gaagaggcct ctttatgcag tcaa 2154

```

&lt;210&gt; 39

&lt;211&gt; 733

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3683905CB1

&lt;400&gt; 39

```

gcgagcgagt tgccgagcgc gccccgtccc tgcgcgcgca tgctcccctg gacggcgctc 60
ggcctggccc tgagcttgcg gctggcgctg gcgcgggagc gcggtccacc 120
agcatcagcc ccccgagggg acctgatgtt cctgctggac agctcagcca gcgtctctca 180
ctacgagttc tcccgggttc gggagtttgt ggggcagctg gtggctccac tgccccctggg 240
caccggggcc ctgctgtgca gtctggtgca cgtgggcagt cggccataca ccgagttccc 300
cttcggccag cacagctcgg gtgaggctgc ccaggatgag gtgctgtctt ctgcccagcg 360
catgggtgac acccacactg gctggcgctt ggtctatgcc aaggaacagc tgtttgtcta 420
agcatcaggt gcccggccag ggggtgccaa agtgcgtgtg tgggtgacag atggcgctc 480
cagcgaccct gtgggcccc ccacgcagga gctcaaggac ctgggcgtca ccgtgttcat 540
tgtcagcacc ggccgaggca acttccctgga gctgtcagcc gctgcctcag cccctgccga 600
gaagcacctg cactttgtgg acgtggatga cctgcacatc attgtccaag agctgagggg 660
ctccattctc gacgcgatgc ggcgcgaagc ttattccctt tagtgagggt taatttttagc 720
ttgcactggc cgt

```

&lt;210&gt; 40

<211> 718  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 4062841CB1

<400> 40  
gccggcgcca gggcaggcgg gcggctggca gctgtggcgc cgacatggct gcgctgggtgg 60  
agccgctggg gctggagcgg gacgtgtccc gggcggttga gctcctcgag cggtccagc 120  
gcagcgggga gctgccgccc cagaagctgc aggcctcca gcgagttctg cagagccgct 180  
tgtgtctcgc tatccgagag gtgtatgagc agctttatga cacgctggac atcaccggca 240  
gcgcccagat ccgagcccat gccacagcca aggccacagt ggctgccttc acagccagcg 300  
agggccacgc acatcccagg gtagtggagc tacccaagac ggatgagggc ctaggcttca 360  
acatcatggg tggcaaagag caaaactcgc ccatctacat ctcccgggtc atcccagggg 420  
gtgtggctga ccgccatgga ggcctcaagc gtggggatca actgttgctg gtgaacgggtg 480  
tgagcgttga gggtagcag catgagaagg cggtaggagc gctgaaggcg gccagggct 540  
cggtagaagc ggttgctcgt tacacaccgc gagtgtgga ggagatggag gcccggttcg 600  
agaagatgcg ctctgcccg cggcgccaac agcatcagag ctactcgctc ttggagtctc 660  
gaggttgaaa ccacagatct ggacgttcac gtgcactctc ttctgtaca gtatttat 718

<210> 41  
<211> 1235  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 6394358CB1

<400> 41  
gccgcgggag ccgcgcgcgc agcgggggag gggagcccc gggggcccc ccaccgccgc 60  
cgctccgcgt cactaccgc gactggatcg gccagagtta ctccgaggtg atgagcctca 120  
acgagcactc catgcaggcg ctgtcctggc gcaagctcta cttgagccgc gccaaactta 180  
aagcctccag ccggacctcg gctctgctct cgggcttcgc catgggtggca atgggtggag 240  
tgcagctgga cgctgaccac gactaccac cggggctgct catcgcttc agtgcttca 300  
ccacagtgtt ggtggctgtg cactgtttg cgtcatgat cagcacctgc atcctgcca 360  
acatcgaggg ggtgagcaac gtgcacaatc tcaactcggg caaggagtcc ccccatgagc 420  
gcatgcaccg ccacatcgag ctggcctggg ccttctccac cgtcatcggc acgtgtctct 480  
tcctagctga ggtggtgctg ctctgctggg tcaagttctt gccctcaag aagcagccag 540  
gccagccaag gccaccagc aagcccccg ccagtggcgc agcagccaac gtcagcaca 600  
gcggcatcac cccgggccag gcagctgcca tcgctcgac caccatcatg gtgccttcg 660  
gctgtacttt tatcgtcttc gccgtccact tctaccgctc actggtttag cataagactg 720  
accgacagtt ccaggagctc aacgagctgg cggagtttgc ccgcttacag gaccagctgg 780  
accacagagg ggaccacccc ctgacgccc gcagccacta tgcttagggc catgtggtct 840  
gggccccttc agtgctttgg ccttacgccc tcccccttga ccttgtcctg cccagcctc 900  
acggacagcc tgccagggg gctgggcttc agcaaggggc agagcgtgga gggaagagga 960  
tttttataag agaaatttct gcactttgaa actgtcctct aagagaataa gcatttctct 1020  
ttcttcagc tccaggtcca cctcctgttg gggggcggtg gggggccaaa gtggggccac 1080  
acaactcgtg tgtcccctct cctcccctgt gccagtcca cctgggtgcc tctcctgtc 1140  
ctgtccgtct caacctccct cccgtccagc attgagtgtg tacatgtgtg tgtgacacat 1200  
aaatatactc ataaggacac ctccaaaaaa aaaaa 1235

<210> 42  
<211> 11648  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 2847752CB1

<400> 42  
atggcgaggc ggccgcgctg gcggggcctc gggggacggt cgacccccat actcctgctc 60  
cttctcctct ctttgttccc cctcagccag gaggagctgg ggggcggtgg gcaccagggc 120  
tgggacccag gcttagctgc cactacgggg ccaagggcgc atatcggtgg cggagcctta 180



gctctttgtc	eggagtcttc	cggggtccgg	gaggatgggg	ggcctggcct	gggggtcagg	240
gagcctatct	tcgtggggct	ccgagggaga	aggcaaaagcg	cccgggaatag	tcgagggccc	300
cctgagcagc	cgaatgagga	gctggggatt	gaacacggcg	tccagccatt	gggcagccgc	360
gaacgagaga	caggacaggg	accaggggtct	gtgttatact	ggcgcccaga	ggtctcctct	420
tgcgggcgga	caggaccttt	gcaaagaggt	agtctgtcac	caggggctct	gtcctcaggg	480
gtcccgggct	cggggaacag	ctcgccccctc	ccttcagact	ttttgattcg	gcaccacggg	540
cccaagccgg	tgctctccca	gcggaacgct	gggacaggct	cccgcaaaag	agtgggcacc	600
gcgcgtgct	gtggggaatt	atgggcaaca	gggagcaagg	gtcagggcga	gagagccacg	660
acatccggag	cagaaaggac	agccccccgg	cggaactgtc	ttccaggggc	ctcgggatct	720
ggccccgagc	tggattcagc	accacgcacg	gcgaggacag	ctcctgcate	aggttcagca	780
ccccgcgagt	ctcgacacagc	tcccagaccg	gcgccccagc	gcatgcgctc	ccggggtctc	840
ttccgctgcc	gcttctctcc	gcagcgcccc	ggggccgcgtc	ccccgggact	ccccggccgt	900
cctgaagcca	ggaaagtaac	ctcggcgaaac	cgggcacgct	ttcgtcgcgc	cgcaaaccgc	960
cacccgcagt	ttccgcagta	caactaccag	acgctgggtg	cggagaatga	ggcagcagge	1020
accgcggtgc	tacgcgtggg	tgctcaggac	ccggacgccc	gcgaggccgg	gcgcctagtc	1080
tactcgctgg	cggcactcat	gaacagccgc	tcgctggagc	tgttcagcat	cgacccgcag	1140
agcggcctta	tcggtacggc	ggcagctctg	gaccgcgaga	gcatggagcg	tcactacctg	1200
cgtgtgaccg	cgcaggacca	cggttcgccc	cgccctctcg	ccaccacgat	ggtggccgtg	1260
acagttagccg	accgcaacga	ccactcgccc	gtttttgagc	aagcgagta	ccgggagacc	1320
cttcgcgaga	atgtggagga	gggctaccct	atcctgcagc	tgctgtccac	tgacggcgac	1380
gcgcccccca	acgccaacct	gcgctaccgc	ttcgtggggc	cgccagctgc	gcgcgctgca	1440
gctgccgccc	ccttcgagat	tgatccacgc	tcgggctca	tcagcaccag	cgggccgagt	1500
gaccgcgagc	acatggaaa	ctatgagctg	gtggtggaag	ccagcgacca	gggcccaggaa	1560
cccgggccgc	gctcggccac	tgtgcgcgta	cacataactg	tgctagacga	gaaccgacaat	1620
gctcctcagt	tcagcgagaa	gcgctacgtg	gcgcaggtgc	gcgaggatgt	gcgcccccac	1680
acagtctgtc	tgcgctcac	ggccactgac	cgggacaagg	acgccaacgg	attgggtgcac	1740
tacaacatca	tcagtggcaa	tagccgtgga	cactttgcca	tcgacagcct	cactggcgag	1800
atccaggtgg	tggcacctct	ggacttcgag	gcagagagag	agtatgcctt	gcgcatacgg	1860
gcgcaggatg	ctggccggcc	accgctgtcc	aacaacacgg	gcctggccag	catccagggtg	1920
gtggacatca	atgaccacat	tcctattttt	gtcagcacgc	ccttccaagt	ttctgtcttg	1980
gaaaatgctc	ccttgggtca	ctcagtcate	cacattcagg	cagtcgatgc	agaccatggg	2040
gagaatgcca	gattggagta	ctccctaact	ggtgtggcac	ctgatactcc	ttttgtgata	2100
aacagcgcca	ctggctgggt	ctctgtgagt	ggtccccctg	accgtgagtc	tgtggagcat	2160
tacttctttg	gtgtggaggc	tcgagaccat	ggctcaccct	cactctctgc	ctcagccagt	2220
gtcacctgta	ctgtgctgga	cgttaatgac	aatcgccctg	agttcacaat	gaaggagtac	2280
cacctacgac	tgaatgagga	tgcagctgtg	ggcaccagtg	tggtcagcgt	gaccgcagta	2340
gacctgtgat	ccaacagtgc	catcagctac	cagatcacag	gcggcaacac	ccggaatcgc	2400
tttgccatca	gcacccaggg	gggtgtgggt	ctggtgactc	tggtctctgc	actggactac	2460
aagcaggaac	gctacttcaa	gctggtacta	actgcactct	accgtgccct	tcattgatcac	2520
tgctatgtgc	acatcaacat	cacagatgcc	aacactcate	ggccgggtctt	tcaaagtgcc	2580
cactactcag	tgaagtgtga	tgaagtgcgg	ccaatgggta	gcaccatagt	ggtcactcagt	2640
gcctctgatg	atgacgtggg	tgagaatgct	cgtatcacct	atctcctgga	ggacaacctg	2700
ccccagttcc	gcattgatgc	agactcagga	gccattacat	tacaggcccc	attagactat	2760
gaggaccagg	tgacctacac	cctggctatc	acagctcggg	acaatggcat	cccacagaag	2820
gcagacacta	cctatgtgga	ggtgatggtc	aatgacgtga	atgacaatgc	tccacaattt	2880
gtggcctccc	actatacagg	gctggctctc	gaggatgccc	cacctttcac	cagtgtcctg	2940
cagatctcag	ccactgaccg	ggatgctcat	gccaatggcc	gggtccagta	cactttccag	3000
aatggtgaag	atggggatgg	agattttacc	attgagccca	cctctggaat	tgctccgtaca	3060
gtaaggcggc	tagaccggga	ggcagtatca	gtgtatgagt	tgactgccta	cgcagtggac	3120
agaggtgtgc	ccccactccg	gactccagtc	agatccagg	tgatggtgca	ggatgtgaac	3180
gacaatgcac	ctgtcttccc	agctgaggag	tttgaggtgc	gggtgaaaaga	gaatagcatt	3240
gtgggctcag	tggtggccca	gatcactgca	gtggaccctg	acgaaggccc	caatgcccac	3300
ataatgtacc	agatcgtgga	ggggaacatc	cctgagctgt	tccaaatgga	catcttctct	3360
ggagaactga	cggcactcat	tgacctagac	tatgaggctc	gccaagaata	tgtgatttgt	3420
gtgcaggcca	catctgctcc	tttggtcagc	cgggccactg	tgacgtcccg	cctggttgac	3480
cagaatgaca	acagccctgt	gctcaacaac	ttccagatcc	tcttcaacaa	ctatgtatcc	3540
aaccgttcag	acaccttccc	gtcgggcatt	attgggcgca	tcccagctta	tgaccccgat	3600
gtctccgacc	acctcttcta	ctcctttgag	cgtggcaatg	agctgcagct	gctggtagtc	3660
aaccagacca	gtggggagct	gcgactcagc	cgaagactag	acaataaccg	cccactgggtg	3720
gcctccatgt	tggtgactgt	cacagatggc	ctgacacagc	tgacggcgca	gtgtgtgtcg	3780
cgcgtgggtca	tcattcacgga	ggagtgtctg	gccaacagcc	tgaccgtgcg	ccttgagaac	3840
atgtggcagg	agcgtctcct	gtcacccgtg	ctgggcccgt	tctctgaggg	cgtggctgcg	3900
gtgctcgcta	cgcccgtgta	ggacgtcttc	atcttcaaca	tccagaacga	cacagacgta	3960
gggggcaccg	tgtctaatgt	gagtttctcg	gcgctagctc	cacgtggggc	cggggcgggc	4020
gctgcagggc	ctcgggttcag	ctcgaggagc	ctgcaagagc	agttgtacgt	gcgcggggcg	4080
gcgctggcgg	ctcgtctcct	gctcgacgta	ctgcccttcg	acgacaacgt	gtgcctgcga	4140
gagccctgtg	agaactacat	gaaatgcgtg	tcggtgctcc	gctttgactc	gtccgcgccc	4200

ttcctggcct	cgccctccac	gctgttccga	cccatccagc	ccatcgctgg	cctgcgctgc	4260
cgctgcccgc	cgggattcac	gggagacttt	tgcgagaccg	agctcgacct	ctgtactctc	4320
aacccatgtc	gcaacggcgg	agcctgcgcg	cggcgcgagg	gaggctacac	gtgcgtctgc	4380
cgcccgcgct	tcaccggaga	ggactgcgag	ctggacaccg	aggccggccg	ctgcgtgccg	4440
ggcgtctgcc	gcaacggggg	cacctgcacc	gacgcgccca	acggcggtct	tcgctgccag	4500
tgcccggcag	gcggcgccct	cgagggcccg	cgctgcgagg	tggtgcgcgc	ctccttcccc	4560
cccagttcgt	tcgtcatgtt	tcgggccttg	cgcgacgat	tcacacctac	gctgtccctc	4620
tcgttcgcga	cagtgcagca	gagcgggctg	ctcttctaca	acgggcgcct	gaacgagaag	4680
cacgacttcc	tggccctgga	actcgtggct	ggccaagtgc	ggctcacata	ttccacgggt	4740
gaatccaaca	ccgtggtcag	ccccacagtt	ccagggggct	tgagtgcagg	gcaatggcat	4800
acagtgcac	tgagatacta	caacaagccc	cggacagatg	ccctaggggg	tgcacagggc	4860
ccctccaagg	acaagggtgg	tgtgctaagc	gtggatgatt	gtgatgtggc	ctgggtctgc	4920
cagtttgggtg	ctgagattgg	caactactca	tgcgcggctg	ctggtgtgca	aacaagctcc	4980
aagaagtccc	tggacctgac	gggcccctct	ctcttggggg	gtgtcccca	cctccccgag	5040
aacttccccg	tatcccataa	ggacttcate	ggctgtatgc	gggacctgca	cattgatggc	5100
cgccgagtgg	acatggcggc	ttttgtcgca	aataatggca	ccatggcagg	ctgccaagcc	5160
aagctacact	tttgtgactc	aggccccctg	acgaacagtg	gcttctgctc	ggacaggtgc	5220
ggcagcttca	gctgcgactg	ccctgtgggc	ttcggcggca	aagactgtca	gcttactatg	5280
gcccattccc	accatttccg	tggcaacggc	acactgagct	ggaactttgg	aagtgcacatg	5340
gctgtgtctg	tgccatggta	cctggggctg	gcatttccga	cacgggcaac	gcagggggtc	5400
ctgatgcaag	tcaggctgg	gccacacagc	acgctccctt	gccagctaga	tcggggggtta	5460
ctgtctgtga	cagtgcaccag	gggctcgggc	cgctgctccc	atctccttct	ggaccaggtg	5520
actgtcagtg	atggccgggtg	gcacgatctg	cggtcggagt	tgcaggagga	accaggtggc	5580
cgccggggcc	accatgtcct	tatggtctca	ctggacttta	gcctcttcca	ggacaccatg	5640
gcgggtggga	gtgagctgca	gggcctgaag	gtaaagcagc	tcacagtggtg	aggcctgccc	5700
cccgagctg	cagaggaggc	tcctcagggt	ctggttggct	gcattccagcc	accgagtga	5760
tgccgacctg	gctgtgttgt	gaccaacgcc	tgtgcctctg	ggccctgccc	acctcacgca	5820
gactgcgggg	acctctggca	gaccttttct	tgcacctgcc	agccaggtta	ctacggccca	5880
ggctgtgtgg	atgcctgcct	cctgaacccc	tgtcagaacc	agggatcatg	ccggcacctg	5940
ccaggagccc	cccatggcta	tacctgtgac	tgtgtgggtg	gctatttccg	gcaccactgt	6000
gagcacagga	tggaccagca	gtgcccacgg	ggctgggtgg	ggagcccaac	ctgtggcccc	6060
tgcaactgtg	atgttcacaa	aggttttgat	cccaactgca	acaagacaaa	tgggcagtgt	6120
cactgcaagg	agttccacta	ccgaccgcgg	ggcagtgact	cttgccctccc	atgtgactgc	6180
tacctgtggg	gctccacctc	gcgctcatgt	gcacccca	gcgggcagtg	ccctgtctgc	6240
ccaggagccc	ttggccgcca	gtgcaacagc	tgtgacagtc	ccttcgcaga	ggtgacagcc	6300
agcggctgcc	gggtgctcta	tgatgcctgc	cctaagtcct	tgagatctgg	tgtgtgggtg	6360
ccccagacaa	agtttggcgt	cctggccaca	gtgccctgtc	cccggggggc	cctgggattg	6420
cggggtgcag	gtgctgctgt	gcggctgtgt	gatgaggccc	agggttggct	ggagcccagc	6480
ctcttcaact	gtacctccc	tgccttctga	gagctcagtc	tgctgctgga	tggcctagag	6540
ctgaacaaga	cggcactgga	tacctggag	gccaagaagc	tggtcagcg	gctacgggtg	6600
gtgactggcc	acactgacca	ctatttttagc	caagatgttc	gagtcactgc	ccgctgtctg	6660
gcccacctgc	tggccttcga	gagccatcag	cagggtctcg	ggctgacagc	cacacaggat	6720
gcccacttca	atgagaatct	gctgtgggccc	ggctctgcac	tgcttgcccc	agagacaggg	6780
gacttgttgg	cggcgctggg	gcagcgggccc	cctgggggct	ccccaggcag	cgcgggactg	6840
gtgaggcacc	tggaggagta	tgcagccaca	ctcgcaagga	atatgggaact	cacatacctg	6900
aatcccatgg	ggctgggtgac	gcctaataatc	atgctcagca	ttgaccgcat	ggagaccccc	6960
agttctcccc	ggggggccccg	tcgctaccct	cgctaccata	gcaacctctt	tcgaggccag	7020
gatgcctggg	atcttcacac	ccatgtgctg	ctgccttccc	agteccccacg	gccatcccca	7080
tctgaagttc	tgcccaaac	cagcagcata	gaaaactcca	ccacctcaag	tgtgtgctcc	7140
ccaccagccc	cgccagagcc	agagcctggg	atctccatta	tcattctcct	cgtttaccgc	7200
accttagggg	gactgtctcc	tgcccagttc	caggcagaac	gccgaggtgc	caggcttctc	7260
cagaaccccc	tcatgaactc	cccgggtggtc	agcgtggctg	tgttccacgg	acgcaacttc	7320
ctaaggggaa	tcctggagtc	ccccatcagc	ctagagtctc	gctgtctaca	gacagcgaat	7380
cggagcaagg	cgatctgtgt	gcagtgggac	ccacctggcc	tggcggagca	gcattggtgtg	7440
tggacagcac	gggactgcga	gctggtgcac	aggaatgggt	cccacgcacg	gtgtcgctgc	7500
agccgggacag	ggacctttgg	ggtcctcatg	gatgcctctc	cccgtagagag	gctggagggc	7560
gacctggagc	tgtggctgtg	gttcacccac	gtggctgtgg	ctgtgtctgt	ggctgcgctg	7620
gtgctgactg	cagccatcct	gctgagcctg	cgcagcctca	agtccaatgt	gcgtgggatc	7680
catgccaatg	tggcagccgc	cctgggggtg	gcagagctcc	tcttctctgt	ggggtattcac	7740
aggacccaca	atcagctggg	gtgcactgca	gtgcacatcc	tcttgcacta	cttcttctctc	7800
agcaccttcc	cgtggctctt	cgtgcagggg	ctgcacctct	accgcatgca	ggttgagcca	7860
cgcaacgtgg	accgcggcgc	catgccttcc	taccatgccc	tgggctgggg	cgctccctgct	7920
gtgctgtctg	gccttgcctg	gggcctggac	cctgagggct	atgggaaccc	tgacttctgc	7980
tggatctcag	tccacgagcc	cctcatctgg	agctttgctg	gccctgttgt	cctgtctcata	8040
gtgatgaacg	ggacctgttt	tctcctcgtc	gcccgcacat	cctgtctccac	agggcagagg	8100
gaggccaaga	agacctctgc	actgaccttt	cgcagctcct	tctgtctgct	tctgtgtgtc	8160
agtgcctcct	ggctctttgg	gctcctggca	gtcaaccaca	gcacctagc	cttccactac	8220

ctccatgctg	gactctgcgg	cctccagggc	ctggcggtgc	tgctgctctt	ctgtgtccta	8280
aatgcagatg	ctcgggctgc	ctggatgcca	gcctgtcttg	gcaggaaggc	agcgcctgag	8340
gaggcaaggc	cagcacctgg	gctgggacct	ggggcctaca	acaacacggc	tctctttgag	8400
gagagtggcc	tcatccgcat	cactctgggc	gcctccaccg	tctcctctgt	gagcagtggc	8460
cgctccggcc	ggaccagga	ccaggacagc	cagcggggcc	gcagctacct	cagggacaat	8520
gtcctgggtc	gacatggctc	agccgctgac	cacactgacc	acagcctcca	ggctcatgct	8580
ggccccactg	acctggacgt	ggccatgttc	catcgagatg	ctggcgacga	ctccgactct	8640
gacagtgcac	tgctccttga	ggaggagagg	agtctctcca	ttccatcttc	agaaagcgag	8700
gacaatggcc	ggacgcgggg	gcgcgtccaa	cggccactct	gccgagcagc	ccagagttag	8760
aggctcctca	cccaccccaa	agatgtggat	ggcaatgacc	tcctgtccta	ctggccagcc	8820
ctgggggagt	gcgaggcagc	cccctgtgct	ctgcagactt	ggggctctga	aaggcgccctg	8880
gggttgagca	ccagcaagga	tgcagctaac	aacaaccagc	cagacccggc	cctgaccagt	8940
gggatcgaga	cttctcttgg	ccgggcccag	cgccagagga	aaggcatcct	gaagaaccgg	9000
ttgcaatacc	cactggtgcc	acagaccgga	ggtgccccctg	agctgtcctg	gtgccgtgca	9060
gccaccttgg	gccaccgtgc	tgtgccagct	gcctctttacg	gtcgcatacta	tgctggcggg	9120
ggcacgggca	gcctttcaca	gccagccagc	cgctactctt	ctagagaaca	gctggacctg	9180
ctcctccggc	ggcaactgag	ccgtgagcga	ctagaggaag	ccccctgccc	tgttctacgt	9240
ccccctgagc	ggccagggtc	ccagggaatgc	atggatgctg	caccaggccg	actggagccc	9300
aaagatcggg	gcagcacctc	gccacggagg	cagccacctc	gggactacct	tggcgccatg	9360
gctggccgct	tggggtcacg	ggatgcgctc	gacttagggg	cacctcgaga	gtggttgagc	9420
acgctgcctc	cgccccggcg	cacccgggac	cttgaccac	agccccccacc	tctgccccctg	9480
tctccccagc	ggcaactctc	aagggaatgc	ctcttgccat	cccgcccgct	ggactctctg	9540
tctaggagct	cgaactctcg	ggagcagctg	gaccaggtgc	ctagccggga	ccccctacga	9600
gaagcccttg	ggccactccc	gcagctgctc	agagctaggg	aggactcggt	cagtggcccc	9660
agccatggcc	cctccacaga	acagttaggac	attctttcct	ccatccttgc	ctctttcaac	9720
tcctcgcccc	tctcctctgt	gcaatcttca	agcacacctc	tgggccccca	caccactgcc	9780
acaccttctg	ccacagcctc	tgtgcttggg	ccctccacgc	cacgttctgc	cacgtctcac	9840
agcatctcgg	agctgtcgcc	agactcagaa	gttcccagaa	gtgagggtca	ctcctgaggg	9900
gatgacggcg	tggacgagga	acagctgagg	gcgacagagg	atctaggcta	acaggagaga	9960
ctccaggagt	gggggcagat	cccaaggcag	cctcctgctc	cccagtgggtg	ggtgccccag	10020
ctctacctgg	tgtggcaggg	ctgaggctcc	atgtgcatct	gtgagcatgc	gtgtgacagg	10080
tgcagagagc	ggggactgga	gggagacttt	tatacgtttt	gtacctttgt	aaccagagag	10140
atgcttatgt	tatttttccag	cttttctgtc	tcctggggggg	tttgaggctg	ggctggggagg	10200
gggagggaga	tagagggaga	gatgcagttt	gaccccat	gggtcctgag	caaaccctat	10260
gctcatctct	ctctccttcc	tggggtggac	tcagatgggt	gggacacatg	ccttccctccc	10320
cctattccac	ccccaaagttg	atctgagtat	cgccaggggc	ccaaagtaca	gaattgttct	10380
ttgcttttta	ttgaatgctc	caaaggccaa	acttctgggg	ctgggggttg	gtcttggaaa	10440
caggggtcct	ctgacttcc	catgggggct	tgctcatacc	gccccctctg	gtggatgtgt	10500
gtgtttatta	tgtggagtcc	ctgccactta	ctgccttatg	acctaggact	gatgtgtgtg	10560
ggtgtgggtg	gagcagctga	tgtcgtgttt	acagagcaag	gcttccctgt	ctccccaggg	10620
gaggggctcg	ggcctctagt	cagacattcc	tgcagagggg	cgggtggagg	gtcattcacc	10680
tgcctctgca	gcaagcaaaa	gttgtctgtg	gtgccatttg	attccctgac	actgccccct	10740
gcttgaattg	attccgaagg	gtaggggtgg	aaggtgagca	aagggagcag	aaacaaggga	10800
attcaagacc	cagaatgtag	gtgccactgc	ctcctatgtt	tacaggatcc	tccgtggccc	10860
taggcacctg	ggctgcagga	agtgaactcc	ttccactcct	cctttattcc	cttaaaaagg	10920
gaaaaatgac	tgttacgacc	ctgttcacaa	aactcttact	tttgctattt	tgtctgctgt	10980
ccagaactga	agactttaaa	attttgttac	tgtttacaag	tccagattca	aaaaatgttt	11040
ttactttgtt	tacaactcaa	aactttgagt	tttacacttt	gtttacagta	gataattttt	11100
tttcctttgt	ttccaagtga	aaggtagggg	aagtgggaga	gggacttgga	ggaccacct	11160
gtgaggacc	tgacctggcc	atcttgaggg	gttttctaac	ccccaggctc	cccaggccga	11220
aggtcagcct	tgagtcccg	ttaacagcag	atccagaaga	ccttgagagt	aggcgtctc	11280
taaccacggg	ggagagtggc	tgtgcagggc	tggggggtgg	tctgtgcaga	cacctctca	11340
cccaccaccc	catgcatact	cttgggaagc	agcttcttgg	gagattagaa	attctacttc	11400
cctgactgga	gctaaatccc	accagccagg	accctaaactc	tccttaccga	gaaggacccc	11460
agctcttgaa	gggctgagt	gcctgctggg	gggtggaggg	tgtctttact	atgtcctagg	11520
tttcgtagat	gccccctctc	ggggttcccc	tcctccagcc	cagcggccct	ctttcctgtc	11580
tgtgtaaat	gttccgtgaa	gccgcgctct	gttttgggaa	taaacttcta	tagaaaacaa	11640
aaaaaaaa						11648

**THIS PAGE BLANK (USPTO)**